

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 August 2001 (23.08.2001)

PCT

(10) International Publication Number
WO 01/60990 A2

(51) International Patent Classification⁷: C12N 9/12,
15/11, 15/63, C07K 16/40, G01N 33/53, C12Q 1/68,
A61K 38/45, 48/00, 39/395, A61P 9/10, 35/00, A01K
67/00

(74) Agent: WRONA, Thomas, J.; Brinks Hofer Gilson & Li-
one, P.O. Box 10087, Chicago, IL 60610 (US).

(21) International Application Number: PCT/US01/04789

(22) International Filing Date: 14 February 2001 (14.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/182,360 14 February 2000 (14.02.2000) US
60/191,261 22 March 2000 (22.03.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants (*for all designated States except US*): CURA-
GEN CORPORATION [US/US]; 322 East Main Street,
Branford, CT 06405 (US). GENENTECH, INC. [US/US];
One DNA Way, South San Francisco, CA 94080 (US).

Published:

— without international search report and to be republished
upon receipt of that report

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): RASTELLI, Luca
[US/US]; 52 Pepperbush Lane, Guilford, CT 06437 (US).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 01/60990 A2

(54) Title: NOVEL SPHINGOSINE KINASES

(57) Abstract: Disclosed herein are novel human nucleic acid sequences that encode polypeptides. Also disclosed are polypeptides encoded by these nucleic acid sequences, and antibodies that immunospecifically-bind to the polypeptide, as well as derivatives, variants, mutants, or fragments of the aforementioned polypeptide, polynucleotide, or antibody. The invention further discloses therapeutic, diagnostic and research methods for diagnosis, treatment, and prevention of disorders involving any one of these novel human nucleic acids and proteins.

The importance of S1P and by consequence the importance of SphK is illustrated by a series of reports that showed their activity in endothelial cell survival and activation by cholesterol via the TNF pathway, in neutrophil activation, in transformed cell motility. ("Tumor necrosis factor- α -mediated signal transduction in human neutrophils: involvement of sphingomyelin metabolites in the priming effect of TNF- α on the fMLP-stimulated superoxide production." Niwa *et al.*, *Life Sci* 2000;66(3):245-56; Overexpression of SphK inhibits chemotactic motility of several transformed cell lines independently of cell surface receptors acting via its substrate, sphingosine-1-phosphate. Wang *et al.*, "Sphingosine-1-phosphate inhibits motility of human breast cancer cells independently of cell surface receptors."

WO 01/60990

PCT/US01/04789

Cancer Res 1999 Dec 15, 59(24):6185-91; "Activation of SphK by tumor necrosis factor- α inhibits apoptosis in human endothelial cells." Xia *et al.*, *J Biol Chem* 1999 Nov 26;274(48):34499-505).

A mouse SphK is described in Kohama *et al.* ("Molecular cloning and functional characterization of murine SphK". *J Biol Chem.* 1998 Sep 11;273(37):23722-8). However, the authors did not have conclusive data on where the mouse protein starts, even if they showed that the mouse cDNAs they obtained have SphK activity *in vivo*. Furthermore, no human sequence has been deposited for this enzyme in GenBank or Geneseq. In their paper, Kohama *et al.* have an alignment that shows only a part of a hypothetical human sequence.

10 SUMMARY OF THE INVENTION

The inventor was able to obtain a full-length human sequence of an SphK with Kozak consensus. Analysis of the full-length human sequence allowed the elucidation of the start of the mouse protein. Furthermore, the inventor discovered a novel SphK that represents a new founder of a subfamily of SphKs.

15 The invention is based in part upon the discovery of novel nucleic acid sequences encoding novel polypeptides. Nucleic acids encoding the polypeptides disclosed in the invention, and derivatives and fragments thereof, will hereinafter be collectively designated as "SphK" nucleic acid or polypeptide sequences.

In one aspect, the invention provides an isolated SphK nucleic acid molecule encoding
20 an SphK polypeptide that includes a nucleic acid sequence that has identity to the nucleic acid disclosed in SEQ ID NO:1, 3, 5, 7, or 9. In some embodiments, the SphK nucleic acid molecule can hybridize under stringent conditions to a nucleic acid sequence complementary to a nucleic acid molecule that includes a protein-coding sequence of an SphK nucleic acid sequence. The invention also includes an isolated nucleic acid that encodes an SphK polypeptide, or a
25 fragment, homolog, analog or derivative thereof. For example, the nucleic acid can encode a polypeptide at least 80% identical to a polypeptide comprising the amino acid sequence of SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15. The nucleic acid can be, for example, a genomic DNA fragment or a cDNA molecule that includes the nucleic acid sequence of any of SEQ ID NOS:1, 3, 5, 7, and 9.

30 Also included in the invention is an oligonucleotide, *e.g.*, an oligonucleotide that includes at least 6 contiguous nucleotides of an SphK nucleic acid (*e.g.*, SEQ ID NO:1, 3, 5, 7, or 9) or a complement of said oligonucleotide.

Also included in the invention are substantially purified SphK polypeptides (SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15). In some embodiments, the SphK polypeptides include an

WO 01/60990

PCT/US01/04789

amino acid sequence that is substantially identical to the amino acid sequence of a human SphK polypeptide.

The invention also features antibodies that immunoselectively-binds to SphK polypeptides.

5 In another aspect, the invention includes pharmaceutical compositions that include therapeutically- or prophylactically-effective amounts of a therapeutic and a pharmaceutically-acceptable carrier. The therapeutic can be, e.g., an SphK nucleic acid, an SphK polypeptide, or an antibody specific for an SphK polypeptide. In a further aspect, the invention includes, in one or more containers, a therapeutically- or prophylactically-effective amount of this
10 pharmaceutical composition.

In a further aspect, the invention includes a method of producing a polypeptide by culturing a cell that includes an SphK nucleic acid, under conditions allowing for expression of the SphK polypeptide encoded by the DNA. If desired, the SphK polypeptide can then be recovered.

15 In another aspect, the invention includes a method of detecting the presence of an SphK polypeptide in a sample. In the method, a sample is contacted with a compound that selectively binds to the polypeptide under conditions allowing for formation of a complex between the polypeptide and the compound. The complex is detected, if present, thereby identifying the SphK polypeptide within the sample.

20 The invention also includes methods to identify specific cell or tissue types based on their expression of an SphK.

Also included in the invention is a method of detecting the presence of an SphK nucleic acid molecule in a sample by contacting the sample with an SphK nucleic acid probe or primer, and detecting whether the nucleic acid probe or primer bound to an SphK nucleic acid molecule
25 in the sample.

In a further aspect, the invention provides a method for modulating the activity of an SphK polypeptide by contacting a cell sample that includes the SphK polypeptide with a compound that binds to the SphK polypeptide in an amount sufficient to modulate the activity of said polypeptide. The compound can be, e.g., a small molecule, such as a nucleic acid, peptide,
30 polypeptide, peptidomimetic, carbohydrate, lipid or other organic (carbon containing) or inorganic molecule, as further described herein.

Also within the scope of the invention is the use of a therapeutic in the manufacture of a medicament for treating or preventing disorders or syndromes including cell proliferative disorders, e.g., cancers, ischemia, or restenosis. The therapeutic can be, e.g., an SphK nucleic

WO 01/60990

PCT/US01/04789

acid, an SphK polypeptide, or an SphK-specific antibody, or biologically-active derivatives or fragments thereof.

The invention further includes a method for screening for a modulator of disorders or syndromes including cell proliferative disorders, *e.g.*, cancers, ischemia, or restenosis. The method includes contacting a test compound with an SphK polypeptide and determining if the test compound binds to said SphK polypeptide. Preferably, the effect of the contacting on enzymatic activity is determined. Compounds that affect SphK enzymatic activity may be modulators of disorders or syndromes.

Also within the scope of the invention is a method for screening for a modulator of activity, or of latency or predisposition to an disorders or syndromes including cell proliferative disorders, *e.g.*, cancers, ischemia, or restenosis, by administering a test compound to a test animal at increased risk for the aforementioned disorders or syndromes. The test animal expresses a recombinant polypeptide encoded by an SphK nucleic acid. Expression or activity of SphK polypeptide is then measured in the test animal, as is expression or activity of the protein in a control animal which recombinantly-expresses SphK polypeptide and is not at increased risk for the disorder or syndrome. Next, the expression of SphK polypeptide in both the test animal and the control animal is compared. A change in the activity of SphK polypeptide in the test animal relative to the control animal indicates the test compound is a modulator of latency of the disorder or syndrome.

In yet another aspect, the invention includes a method for determining the presence of or predisposition to a disease associated with altered levels of an SphK polypeptide, an SphK nucleic acid, or both, in a subject (*e.g.*, a human subject). The method includes measuring the amount of the SphK polypeptide in a test sample from the subject and comparing the amount of the polypeptide in the test sample to the amount of the SphK polypeptide present in a control sample. An alteration in the level of the SphK polypeptide in the test sample as compared to the control sample indicates the presence of or predisposition to a disease in the subject.

Alternatively, rather than measuring the amount of SphK polypeptide, SphK enzymatic activity can be measured. Preferably, the predisposition includes cell proliferative disorders, *e.g.*, cancers, ischemia, or restenosis. Also, the expression levels of the new polypeptides of the invention can be used in a method to screen for various cancers.

In a further aspect, the invention includes a method of treating or preventing a pathological condition associated with a disorder in a mammal by administering to the subject an SphK polypeptide, an SphK nucleic acid, or an SphK-specific antibody to a subject (*e.g.*, a human subject), in an amount sufficient to alleviate or prevent the pathological condition. In

WO 01/60990

PCT/US01/04789

preferred embodiments, the disorder includes cell proliferative disorders, e.g., cancers, ischemia, or restenosis.

In yet another aspect, the invention can be used in a method to identity the cellular receptors and downstream effectors of the invention by any one of a number of techniques commonly employed in the art. These include but are not limited to the two-hybrid system, affinity purification, co-precipitation with antibodies or other specific-interacting molecules.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

15

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. Alignment of the novel human SphK (SPkinaseinHSDA59H18; SEQ ID NO:6) with the amino acid sequence of CAB62977 (SEQ ID. NO:11).

FIG. 2. Alignment of the novel human SphK (SPkinaseinHSDA59H18; SEQ ID NO:6) with the rat (AI237625; SEQ ID NO:8) and mouse (52180312; SEQ ID NO:10) orthologs.

FIG. 3. Alignment of the novel human SphK (SPkinaseinHSDA59H18; SEQ ID NO:6) with the human SphK of Example 1 (coorrected_human_sphingosine; SEQ ID NO:2), its mouse homolog (AF068748_EXT-2; SEQ ID NO:4), along with homologs from other species (Q06147 (*Saccharomyces ceravisiae*; SEQ ID NO:12); Q12246 (*Saccharomyces ceravisiae*; SEQ ID NO:13); O14159 (*Schizosaccharomyces pombe*; SEQ ID NO:14); and Q18425(*Caenorhabditis elegans*; SEQ ID NO:15)).

FIG. 4. Protein distance analysis of the SphKs.

FIG. 5. Alignment of SPkinaseinHSDA59H18 (SEQ ID NO:6) and 80432911.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

30

DETAILED DESCRIPTION

The invention is based, in part, upon the discovery of novel nucleic acid sequences that encode novel polypeptides, particularly sphingosine kinases. The nucleic acids, and their encoded polypeptides, are collectively designated herein as "SphK".

WO 01/60990

PCT/US01/04789

The novel SphK nucleic acids of the invention include the nucleic acids whose sequences are provided as SEQ ID NO:1, 3, 5, 7, or 9, or a fragment thereof. The invention also includes a mutant or variant SphK nucleic acid, any of whose bases may be changed from the corresponding base listed in SEQ ID NO:1, 3, 5, 7, or 9 while still encoding a protein that
 5 maintains the activities and physiological functions of the SphK protein fragment, or a fragment of such a nucleic acid. The invention further includes nucleic acids whose sequences are complementary to those just described, including complementary nucleic acid fragments. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications. Such modifications include, by way of
 10 nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject. In the mutant or variant nucleic acids, and their complements, up to 20% or more of the bases may be so changed.

15 The novel SphK proteins of the invention include the protein fragments whose sequences are provided as SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15. The invention also includes an SphK mutant or variant protein, any of whose residues may be changed from the corresponding residue listed in SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15 while still encoding a protein that maintains its native activities and physiological functions, or a functional fragment thereof. In
 20 the mutant or variant SphK protein, up to 20% or more of the residues may be so changed. The invention further encompasses antibodies and antibody fragments, such as F_{ab} or (F_{ab})₂, that bind immunospecifically to any of the SphK proteins of the invention.

The SphK nucleic acids and proteins of the invention are useful in potential therapeutic applications implicated in various cancers, tumors and similar neoplastic diseases. For example,
 25 a cDNA encoding a polypeptide of SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15 may be useful in gene therapy, and the polypeptide itself may be useful when administered to a subject in need thereof. The novel nucleic acids or polypeptides of the present, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed. These materials are further useful in the generation of
 30 antibodies that bind immunospecifically to the novel substances of the invention for use in therapeutic or diagnostic methods.

SphK Nucleic Acids and Polypeptides

One aspect of the invention pertains to isolated nucleic acid molecules that encode SphK polypeptides or biologically-active portions thereof. Also included in the invention are nucleic

WO 01/60990

PCT/US01/04789

acid fragments sufficient for use as hybridization probes to identify SphK-encoding nucleic acids (e.g., SphK mRNAs) and fragments for use as PCR primers for the amplification and/or mutation of SphK nucleic acid molecules. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (e.g., cDNA or genomic DNA), RNA molecules (e.g., mRNA), analogs of the DNA or RNA generated using nucleotide analogs, and derivatives, fragments and homologs thereof. The nucleic acid molecule may be single-stranded or double-stranded, but preferably is comprised double-stranded DNA.

An SphK nucleic acid can encode a mature SphK polypeptide. As used herein, a "mature" form of a polypeptide or protein disclosed in the present invention is the product of a naturally occurring polypeptide or precursor form or proprotein. The naturally occurring polypeptide, precursor or proprotein includes, by way of nonlimiting example, the full length gene product, encoded by the corresponding gene. Alternatively, it may be defined as the polypeptide, precursor or proprotein encoded by an open reading frame described herein. The product "mature" form arises, again by way of nonlimiting example, as a result of one or more naturally occurring processing steps as they may take place within the cell, or host cell, in which the gene product arises. Examples of such processing steps leading to a "mature" form of a polypeptide or protein include the cleavage of the N-terminal methionine residue encoded by the initiation codon of an open reading frame, or the proteolytic cleavage of a signal peptide or leader sequence. Thus a mature form arising from a precursor polypeptide or protein that has residues 1 to N, where residue 1 is the N-terminal methionine, would have residues 2 through N remaining after removal of the N-terminal methionine. Alternatively, a mature form arising from a precursor polypeptide or protein having residues 1 to N, in which an N-terminal signal sequence from residue 1 to residue M is cleaved, would have the residues from residue M+1 to residue N remaining. Further as used herein, a "mature" form of a polypeptide or protein may arise from a step of post-translational modification other than a proteolytic cleavage event. Such additional processes include, by way of non-limiting example, glycosylation, myristoylation or phosphorylation. In general, a mature polypeptide or protein may result from the operation of only one of these processes, or a combination of any of them.

The term "probes", as utilized herein, refers to nucleic acid sequences of variable length, preferably between at least about 10 nucleotides (nt), 100 nt, or as many as approximately, e.g., 6,000 nt, depending upon the specific use. Probes are used in the detection of identical, similar, or complementary nucleic acid sequences. Longer length probes are generally obtained from a natural or recombinant source, are highly specific, and much slower to hybridize than shorter-length oligomer probes. Probes may be single- or double-stranded and designed to have

WO 01/60990

PCT/US01/04789

specificity in PCR, membrane-based hybridization technologies, array technologies, or ELISA-like technologies.

The term "isolated" nucleic acid molecule, as utilized herein, is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences that naturally flank the nucleic acid (i.e., sequences located at the 5'- and 3'-termini of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated SphK nucleic acid molecules can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell/tissue from which the nucleic acid is derived (e.g., brain, heart, liver, spleen, etc.). Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material or culture medium when produced by recombinant techniques, or of chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the invention, e.g., a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1, 3, 5, 7, or 9, or a complement of this aforementioned nucleotide sequence, can be isolated using standard molecular biology techniques and the sequence information provided herein. Using all or a portion of the nucleic acid sequence of SEQ ID NO:1, 3, 5, 7, or 9 as a hybridization probe, SphK molecules can be isolated using standard hybridization and cloning techniques (e.g., as described in Sambrook, *et al.*, (eds.), MOLECULAR CLONING: A LABORATORY MANUAL 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989; and Ausubel, *et al.*, (eds.), CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, NY, 1993.)

A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to SphK nucleotide sequences can be prepared by standard synthetic techniques, e.g., using an automated DNA synthesizer.

As used herein, the term "oligonucleotide" refers to a series of linked nucleotide residues, which oligonucleotide has a sufficient number of nucleotide bases to be used in a PCR reaction. A short oligonucleotide sequence may be based on, or designed from, a genomic or cDNA sequence and is used to amplify, confirm, or reveal the presence of an identical, similar or complementary DNA or RNA in a particular cell or tissue. Oligonucleotides comprise portions of a nucleic acid sequence having about 10 nt, 50 nt, or 100 nt in length, preferably about 15 nt to 30 nt in length. In one embodiment of the invention, an oligonucleotide

WO 01/60990

PCT/US01/04789

comprising a nucleic acid molecule less than 100 nt in length would further comprise at least 6 contiguous nucleotides of SEQ ID NO:1, 3, 5, 7, or 9, or a complement thereof.

Oligonucleotides may be chemically synthesized and may also be used as probes.

In another embodiment, an isolated nucleic acid molecule of the invention comprises a
 5 nucleic acid molecule that is a complement of the nucleotide sequence shown in SEQ ID NO:1, 3, 5, 7, or 9, or a portion of this nucleotide sequence (e.g., a fragment that can be used as a probe or primer or a fragment encoding a biologically-active portion of an SphK polypeptide). A nucleic acid molecule that is complementary to the nucleotide sequence shown in SEQ ID NO:1, 3, 5, 7, or 9, is one that is sufficiently complementary to the nucleotide sequence shown in SEQ
 10 ID NO:1, 3, 5, 7, or 9, that it can hydrogen bond with little or no mismatches to the nucleotide sequence shown in SEQ ID NO:1, 3, 5, 7, or 9, thereby forming a stable duplex.

As used herein, the term "complementary" refers to Watson-Crick or Hoogsteen base pairing between nucleotides units of a nucleic acid molecule, and the term "binding" means the physical or chemical interaction between two polypeptides or compounds or associated
 15 polypeptides or compounds or combinations thereof. Binding includes ionic, non-ionic, van der Waals, hydrophobic interactions, and the like. A physical interaction can be either direct or indirect. Indirect interactions may be through or due to the effects of another polypeptide or compound. Direct binding refers to interactions that do not take place through, or due to, the effect of another polypeptide or compound, but instead are without other substantial chemical
 20 intermediates.

Fragments provided herein are defined as sequences of at least 6 (contiguous) nucleic acids or at least 4 (contiguous) amino acids, a length sufficient to allow for specific hybridization in the case of nucleic acids or for specific recognition of an epitope in the case of amino acids, respectively, and are at most some portion less than a full length sequence.
 25 Fragments may be derived from any contiguous portion of a nucleic acid or amino acid sequence of choice. Derivatives are nucleic acid sequences or amino acid sequences formed from the native compounds either directly or by modification or partial substitution. Analogs are nucleic acid sequences or amino acid sequences that have a structure similar to, but not identical to, the native compound but differ from it in respect to certain components or side chains. Analogs
 30 may be synthetic or from a different evolutionary origin and may have a similar or opposite metabolic activity compared to wild type. Homologs are nucleic acid sequences or amino acid sequences of a particular gene that are derived from different species.

Derivatives and analogs may be full length or other than full length, if the derivative or analog contains a modified nucleic acid or amino acid, as described below. Derivatives or
 35 analogs of the nucleic acids or proteins of the invention include, but are not limited to,

WO 01/60990

PCT/US01/04789

molecules comprising regions that are substantially homologous to the nucleic acids or proteins of the invention, in various embodiments, by at least about 70%, 80%, or 95% identity (with a preferred identity of 80-95%) over a nucleic acid or amino acid sequence of identical size or when compared to an aligned sequence in which the alignment is done by a computer homology program known in the art, or whose encoding nucleic acid is capable of hybridizing to the complement of a sequence encoding the aforementioned proteins under stringent, moderately stringent, or low stringent conditions. See *e.g.* Ausubel, *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, NY, 1993, and below.

A "homologous nucleic acid sequence" or "homologous amino acid sequence," or variations thereof, refer to sequences characterized by a homology at the nucleotide level or amino acid level as discussed above. Homologous nucleotide sequences encode those sequences coding for isoforms of SphK polypeptides. Isoforms can be expressed in different tissues of the same organism as a result of, for example, alternative splicing of RNA. Alternatively, isoforms may be encoded by different genes. In the invention, homologous nucleotide sequences include nucleotide sequences encoding for an SphK polypeptide of species other than humans, including, but not limited to: vertebrates, and thus can include, *e.g.*, frog, mouse, rat, rabbit, dog, cat, cow, horse, and other organisms. Homologous nucleotide sequences also include, but are not limited to, naturally occurring allelic variations and mutations of the nucleotide sequences set forth herein. A homologous nucleotide sequence does not, however, include the exact nucleotide sequence encoding human SphK protein. Homologous nucleic acid sequences include those nucleic acid sequences that encode conservative amino acid substitutions (see below) in SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15, as well as a polypeptide possessing SphK biological activity. Various biological activities of the SphK proteins are described below.

An SphK polypeptide is encoded by the open reading frame ("ORF") of an SphK nucleic acid. An ORF corresponds to a nucleotide sequence that could potentially be translated into a polypeptide. A stretch of nucleic acids comprising an ORF is uninterrupted by a stop codon. An ORF that represents the coding sequence for a full protein begins with an ATG "start" codon and terminates with one of the three "stop" codons, namely, TAA, TAG, or TGA. For the purposes of this invention, an ORF may be any part of a coding sequence, with or without a start codon, a stop codon, or both. For an ORF to be considered as a good candidate for coding for a *bona fide* cellular protein, a minimum size requirement is often set, *e.g.*, a stretch of DNA that would encode a protein of 50 amino acids or more.

The nucleotide sequences determined from the SphK genes allows for the generation of probes and primers designed for use in identifying and/or cloning SphK homologues in other cell types, *e.g.* from other tissues, as well as SphK homologues from other vertebrates. The

WO 01/60990

PCT/US01/04789

probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 25, 50, 100, 150, 200, 250, 300, 350 or 400 consecutive sense strand nucleotide sequence of SEQ ID NO:1, 3, 5, 7, or 9; or an anti-sense strand nucleotide sequence of SEQ ID NO:1, 3, 5, 7, or 9; or of a naturally occurring mutant of SEQ ID NO:1, 3, 5, 7, or 9.

Probes based on the human SphK nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In various embodiments, the probe further comprises a label group attached thereto, e.g. the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissues which mis-express an SphK protein, such as by measuring a level of an SphK-encoding nucleic acid in a sample of cells from a subject e.g., detecting SphK mRNA levels or determining whether a genomic SphK gene has been mutated or deleted.

"A polypeptide having a biologically-active portion of an SphK polypeptide" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. A nucleic acid fragment encoding a "biologically-active portion of SphK" can be prepared by isolating a portion of SEQ ID NO:1, 3, 5, 7, or 9, that encodes a polypeptide having an SphK biological activity (the biological activities of the SphK proteins are described below), expressing the encoded portion of SphK protein (e.g., by recombinant expression *in vitro*) and assessing the activity of the encoded portion of SphK.

SphK Nucleic Acid and Polypeptide Variants

The invention further encompasses nucleic acid molecules that differ from the nucleotide sequence shown in SEQ ID NO:1, 3, 5, 7, or 9, due to degeneracy of the genetic code and thus encode the same SphK protein as that encoded by the nucleotide sequence shown in SEQ ID NO:1, 3, 5, 7, or 9. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15.

In addition to the SphK nucleotide sequences shown in SEQ ID NOS:1, 3, 5, 7, and 9, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of the SphK polypeptides may exist within a population (e.g., the human population). Such genetic polymorphism in the SphK genes may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame

WO 01/60990

PCT/US01/04789

(ORF) encoding an SphK protein, preferably a vertebrate SphK protein. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of the SphK genes. Any and all such nucleotide variations and resulting amino acid polymorphisms in the SphK polypeptides, which are the result of natural allelic variation and that do not alter the functional activity of the SphK polypeptides, are intended to be within the scope of the invention.

Moreover, nucleic acid molecules encoding SphK proteins from additional species, and thus that have a nucleotide sequence that differs from the sequence of SEQ ID NO:1, 3, 5, 7, or 9, are intended to be within the scope of the invention. Nucleic acid molecules corresponding to natural allelic variants and homologues of the SphK cDNAs of the invention can be isolated based on their homology to the human SphK nucleic acids disclosed herein using the human cDNAs, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 6 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 3, 5, 7, or 9. In another embodiment, the nucleic acid is at least 10, 25, 50, 100, 250, 500, 750, 1000, 1500, or 2000 or more nucleotides in length. In yet another embodiment, an isolated nucleic acid molecule of the invention hybridizes to the coding region. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% identical to each other typically remain hybridized to each other.

Homologs (*i.e.*, nucleic acids encoding SphK proteins derived from species other than that from which the nucleic acid was originally isolated) or other related sequences (*e.g.*, paralogs) can be obtained by low, moderate or high stringency hybridization with all or a portion of the particular sequence as a probe using methods well known in the art for nucleic acid hybridization and cloning.

As used herein, the phrase "stringent hybridization conditions" refers to conditions under which a probe, primer or oligonucleotide will hybridize to its target sequence, but preferably to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures than shorter sequences. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the probes complementary to the target sequence hybridize to the target sequence at equilibrium. Since the target sequences are generally present at excess, at T_m , 50% of the

WO 01/60990

PCT/US01/04789

probes are occupied at equilibrium. Typically, stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes, primers or oligonucleotides (*e.g.*, 10 nt to 50 nt) and at least about 60°C for longer probes, primers and oligonucleotides. Stringent conditions may also be achieved with the addition of destabilizing agents, such as formamide.

Stringent conditions are known to those skilled in the art and can be found in Ausubel, *et al.*, (eds.), CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Preferably, the conditions are such that sequences at least about 65%, 70%, 75%, 85%, 90%, 95%, 98%, or 99% identical to each other typically remain hybridized to each other. A non-limiting example of stringent hybridization conditions are hybridization in a high salt buffer comprising 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 mg/ml denatured salmon sperm DNA at 65°C, followed by one or more washes in 0.2X SSC, 0.01% BSA at 50°C. An isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence SEQ ID NO:1, 3, 5, 7, or 9 may corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

In a second embodiment, a nucleic acid sequence that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 3, 5, 7, or 9, or fragments, analogs or derivatives thereof, under conditions of moderate stringency is provided. A non-limiting example of moderate stringency hybridization conditions are hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100 mg/ml denatured salmon sperm DNA at 55°C, followed by one or more washes in 1X SSC, 0.1% SDS at 37°C. Other conditions of moderate stringency that may be used are well-known within the art. See, *e.g.*, Ausubel, *et al.* (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, NY, and Kriegler, 1990; GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY.

In a third embodiment, a nucleic acid that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 3, 5, 7, or 9, or fragments, analogs or derivatives thereof, under conditions of low stringency, is provided. A non-limiting example of low stringency hybridization conditions are hybridization in 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 mg/ml denatured salmon sperm DNA, 10% (wt/vol) dextran sulfate at 40°C, followed by one or more washes in 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS at 50°C. Other conditions of low stringency that may be used are well known in the art (*e.g.*, as employed for cross-species

WO 01/60990

PCT/US01/04789

hybridizations). See, e.g., Ausubel, *et al.* (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, NY, and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY; Shilo and Weinberg, 1981. *Proc Natl Acad Sci USA* 78: 6789-6792.

5 **Conservative Mutations**

In addition to naturally-occurring allelic variants of SphK sequences that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequence of SEQ ID NO:1, 3, 5, 7, or 9, thereby leading to changes in the amino acid sequences of the encoded SphK proteins, without altering the functional ability of said SphK proteins. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in the sequence of SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequences of the SphK proteins without altering their biological activity, whereas an "essential" amino acid residue is required for such biological activity. For example, amino acid residues that are conserved among the SphK proteins of the invention are predicted to be particularly non-amenable to alteration. Amino acids for which conservative substitutions can be made are well-known within the art.

Another aspect of the invention pertains to nucleic acid molecules encoding SphK proteins that contain changes in amino acid residues that are not essential for activity. Such SphK proteins differ in amino acid sequence from SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15, yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 45% identical to the amino acid sequences of SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15. Preferably, the protein encoded by the nucleic acid molecule is at least about 60% identical to SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15; more preferably at least about 70% identical to SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15; still more preferably at least about 80% identical to SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15; even more preferably at least about 90% identical to SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15; and most preferably at least about 95% identical to SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15.

30 An isolated nucleic acid molecule encoding an SphK protein homologous to the protein of SEQ ID NOS:2, 4, 6, 8, 10, 12, 13, 14, or 15, can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO:1, 3, 5, 7, or 9, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein.

30 In yet another embodiment, a mutant SphK protein can be assayed for the ability to regulate a specific biological function (e.g., regulation of an S1P responsive signal transduction pathway).

WO 01/60990

PCT/US01/04789

Antisense Nucleic Acids

Another aspect of the invention pertains to isolated antisense nucleic acid molecules that are hybridizable to or complementary to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 3, 5, 7, or 9, or fragments, analogs or derivatives thereof. An
 5 "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein (e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence). In specific aspects, antisense nucleic acid molecules are provided that comprise a sequence complementary to at least about
 10 10, 25, 50, 100, 250 or 500 nucleotides or an entire SphK coding strand, or to only a portion thereof. Nucleic acid molecules encoding fragments, homologs, derivatives and analogs of an SphK protein of SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15; or antisense nucleic acids complementary to an SphK nucleic acid sequence of SEQ ID NO:1, 3, 5, 7, or 9, are additionally provided.

In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region"
 15 of the coding strand of a nucleotide sequence encoding an SphK protein. The term "coding region" refers to the region of the nucleotide sequence comprising codons that are translated into amino acid residues. In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding the SphK protein. The term "noncoding region" refers to 5' and 3' sequences that flank the coding region that are
 20 not translated into amino acids (i.e., also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences encoding the SphK proteins disclosed herein, antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of SphK mRNA, but more preferably is an oligonucleotide that is
 25 antisense to only a portion of the coding or noncoding region of SphK mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of SphK mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis or enzymatic ligation reactions using procedures known in
 30 the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally-occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids (e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used).

WO 01/60990

PCT/US01/04789

Examples of modified nucleotides that can be used to generate the antisense nucleic acid include: 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding an SphK protein to thereby inhibit expression of the protein (*e.g.*, by inhibiting transcription and/or translation). The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site by syringe, stent, or by catheter. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface (*e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens). The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient nucleic acid molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands

WO 01/60990

PCT/US01/04789

run parallel to each other. See, e.g., Gaultier, *et al.*, 1987. *Nucl. Acids Res.* 15: 6625-6641. The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (see, e.g., Inoue, *et al.* 1987. *Nucl. Acids Res.* 15: 6131-6148) or a chimeric RNA-DNA analogue (see, e.g., Inoue, *et al.*, 1987. *FEBS Lett.* 215: 327-330.

5 **Ribozymes and PNA Moieties**

Nucleic acid modifications include, by way of non-limiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in
10 therapeutic applications in a subject.

In one embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity that are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (e.g., hammerhead ribozymes as described in Haselhoff and Gerlach 1988. *Nature*
15 334: 585-591) can be used to catalytically cleave SphK mRNA transcripts to thereby inhibit translation of SphK mRNA. A ribozyme having specificity for an SphK-encoding nucleic acid can be designed based upon the nucleotide sequence of an SphK cDNA disclosed herein (*i.e.*, SEQ ID NO:1, 3, 5, 7, or 9). For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the
20 nucleotide sequence to be cleaved in an SphK-encoding mRNA. See, e.g., U.S. Patent 4,987,071 to Cech, *et al.* and U.S. Patent 5,116,742 to Cech, *et al.* SphK mRNA can also be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, e.g., Bartel *et al.*, (1993) *Science* 261:1411-1418.

Alternatively, SphK gene expression can be inhibited by targeting nucleotide sequences
25 complementary to the regulatory region of the SphK nucleic acid (e.g., the SphK promoter and/or enhancers) to form triple helical structures that prevent transcription of the SphK gene in target cells. See, e.g., Helene, 1991. *Anticancer Drug Des.* 6: 569-84; Helene, *et al.* 1992. *Ann. N.Y. Acad. Sci.* 660: 27-36; Maher, 1992. *Bioassays* 14: 807-15.

In various embodiments, the SphK nucleic acids can be modified at the base moiety,
30 sugar moiety or phosphate backbone to improve, e.g., the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids. See, e.g., Hyrup, *et al.*, 1996. *Bioorg Med Chem* 4: 5-23. As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics (e.g., DNA mimics) in which the deoxyribose phosphate backbone is replaced by a
35 pseudopeptide backbone and only the four natural nucleobases are retained. The neutral

WO 01/60990

PCT/US01/04789

backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup, *et al.*, 1996. *supra*; Perry-O'Keefe, *et al.*, 1996. *Proc. Natl. Acad. Sci. USA* 93: 14670-14675.

5 PNAs of SphK can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, *e.g.*, inducing transcription or translation arrest or inhibiting replication. PNAs of SphK can also be used, for example, in the analysis of single base pair mutations in a gene (*e.g.*, PNA directed PCR clamping; as artificial restriction enzymes when used in combination with
10 other enzymes, *e.g.*, S_I nucleases (*see*, Hyrup, *et al.*, 1996. *supra*); or as probes or primers for DNA sequence and hybridization (*see*, Hyrup, *et al.*, 1996, *supra*; Perry-O'Keefe, *et al.*, 1996. *supra*).

In another embodiment, PNAs of SphK can be modified, *e.g.*, to enhance their stability or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of
15 PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras of SphK can be generated that may combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes (*e.g.*, RNase H and DNA polymerases) to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using
20 linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (*see*, Hyrup, *et al.*, 1996. *supra*). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup, *et al.*, 1996. *supra* and Finn, *et al.*, 1996. *Nucl Acids Res* 24: 3357-3363. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry, and modified nucleoside analogs, *e.g.*,
25 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used between the PNA and the 5' end of DNA. *See, e.g.*, Mag, *et al.*, 1989. *Nucl Acid Res* 17: 5973-5988. PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment. *See, e.g.*, Finn, *et al.*, 1996. *supra*. Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment. *See, e.g.*,
30 Petersen, *et al.*, 1975. *Bioorg. Med. Chem. Lett.* 5: 1119-11124.

In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (*see, e.g.*, Letsinger, *et al.*, 1989. *Proc. Natl. Acad. Sci. U.S.A.* 86: 6553-6556; Lemaitre, *et al.*, 1987. *Proc. Natl. Acad. Sci.* 84: 648-652; PCT Publication No. WO88/09810)
35 or the blood-brain barrier (*see, e.g.*, PCT Publication No. WO 89/10134). In addition,

WO 01/60990

PCT/US01/04789

oligonucleotides can be modified with hybridization triggered cleavage agents (*see, e.g., Krol, et al., 1988. BioTechniques 6:958-976*) or intercalating agents (*see, e.g., Zon, 1988. Pharm. Res. 5: 539-549*). To this end, the oligonucleotide may be conjugated to another molecule, *e.g., a*
peptide, a hybridization triggered cross-linking agent, a transport agent, a
5 hybridization-triggered cleavage agent, and the like.

SphK Polypeptides

A polypeptide according to the invention includes a polypeptide including the amino acid sequence of SphK polypeptides whose sequences are provided in SEQ ID NOS:2, 4, 6, 8, 10, 12, 13, 14, and 15. The invention also includes a mutant or variant protein any of whose
10 residues may be changed from the corresponding residues shown in SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15, while still encoding a protein that maintains its SphK activities and physiological functions, or a functional fragment thereof.

In general, an SphK variant that preserves SphK-like function includes any variant in which residues at a particular position in the sequence have been substituted by other amino
15 acids, and further include the possibility of inserting an additional residue or residues between two residues of the parent protein as well as the possibility of deleting one or more residues from the parent sequence. Any amino acid substitution, insertion, or deletion is encompassed by the invention. In favorable circumstances, the substitution is a conservative substitution as defined above.

20 One aspect of the invention pertains to isolated SphK proteins, and biologically-active portions thereof, or derivatives, fragments, analogs or homologs thereof. Also provided are polypeptide fragments suitable for use as immunogens to raise anti-SphK antibodies. In one embodiment, native SphK proteins can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment,
25 SphK proteins are produced by recombinant DNA techniques. Alternative to recombinant expression, an SphK protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" polypeptide or protein or biologically-active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue
30 source from which the SphK protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of SphK proteins in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly-produced. In one embodiment, the language "substantially free of cellular material" includes preparations of SphK

WO 01/60990

PCT/US01/04789

proteins having less than about 30% (by dry weight) of non-SphK proteins (also referred to herein as a "contaminating protein"), more preferably less than about 20% of nonSphK proteins, still more preferably less than about 10% of non-SphK proteins, and most preferably less than about 5% of non-SphK proteins. When the SphK protein or biologically-active portion thereof is recombinantly-produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the SphK protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of SphK proteins in which the protein is separated from chemical precursors or other chemicals that are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of SphK proteins having less than about 30% (by dry weight) of chemical precursors or non-SphK chemicals, more preferably less than about 20% chemical precursors or non-SphK chemicals, still more preferably less than about 10% chemical precursors or non-SphK chemicals, and most preferably less than about 5% chemical precursors or non-SphK chemicals.

Biologically-active portions of SphK proteins include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequences of the SphK proteins (*e.g.*, the amino acid sequence shown in SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15) that include fewer amino acids than the full-length SphK proteins, and exhibit at least one activity of an SphK protein. Typically, biologically-active portions comprise a domain or motif with at least one activity of the SphK protein. A biologically-active portion of an SphK protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acid residues in length.

Moreover, other biologically-active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native SphK protein.

In an embodiment, the SphK protein has an amino acid sequence shown in SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15. In other embodiments, the SphK protein is substantially homologous to SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15, and retains the functional activity of the protein of SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail, below. Accordingly, in another embodiment, the SphK protein is a protein that comprises an amino acid sequence at least about 45% homologous to the amino acid sequence of SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15, and retains the functional activity of the SphK proteins of SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15.

WO 01/60990

PCT/US01/04789

Determining Homology Between Two or More Sequences

To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are homologous at that position (i.e., as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity").

The nucleic acid sequence homology may be determined as the degree of identity between two sequences. The homology may be determined using computer programs known in the art, such as GAP software provided in the GCG program package. See, Needleman and Wunsch, 1970. *J Mol Biol* 48: 443-453. Using GCG GAP software with the following settings for nucleic acid sequence comparison: GAP creation penalty of 5.0 and GAP extension penalty of 0.3, the coding region of the analogous nucleic acid sequences referred to above exhibits a degree of identity preferably of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99%, with the CDS (encoding) part of the DNA sequence shown in SEQ ID NO:1, 3, 5, 7, or 9.

The term "sequence identity" refers to the degree to which two polynucleotide or polypeptide sequences are identical on a residue-by-residue basis over a particular region of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over that region of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, U, or I, in the case of nucleic acids) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the region of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The term "substantial identity" as used herein denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 80 percent sequence identity, preferably at least 85 percent identity and often 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison region.

Chimeric and Fusion Proteins

The invention also provides SphK chimeric or fusion proteins. As used herein, an SphK "chimeric protein" or "fusion protein" comprises an SphK polypeptide operatively-linked to a non-SphK polypeptide. A "SphK polypeptide" refers to a polypeptide having an amino acid sequence corresponding to an SphK protein (SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15),

WO 01/60990

PCT/US01/04789

whereas a "non-SphK polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein that is not substantially homologous to the SphK protein, *e.g.*, a protein that is different from the SphK protein and that is derived from the same or a different organism. Within an SphK fusion protein the SphK polypeptide can correspond to all or a portion of an SphK protein. In one embodiment, an SphK fusion protein comprises at least one biologically-active portion of an SphK protein. In another embodiment, an SphK fusion protein comprises at least two biologically-active portions of an SphK protein. In yet another embodiment, an SphK fusion protein comprises at least three biologically-active portions of an SphK protein. Within the fusion protein, the term "operatively-linked" is intended to indicate that the SphK polypeptide and the non-SphK polypeptide are fused in-frame with one another. The non-SphK polypeptide can be fused to the N-terminus or C-terminus of the SphK polypeptide.

In one embodiment, the fusion protein is a GST-SphK fusion protein in which the SphK sequences are fused to the C-terminus of the GST (glutathione S-transferase) sequences. Such fusion proteins can facilitate the purification of recombinant SphK polypeptides.

In another embodiment, the fusion protein is an SphK protein containing a heterologous signal sequence at its N-terminus. In certain host cells (*e.g.*, mammalian host cells), expression and/or secretion of SphK can be increased through use of a heterologous signal sequence.

In yet another embodiment, the fusion protein is an SphK-immunoglobulin fusion protein in which the SphK sequences are fused to sequences derived from a member of the immunoglobulin protein family. The SphK-immunoglobulin fusion proteins can be used to affect the bioavailability of an SphK substrate. Inhibition of the SphK substrate/ SphK interaction may be useful therapeutically for both the treatment of proliferative and differentiative disorders, as well as modulating (*e.g.* promoting or inhibiting) cell survival. Moreover, the SphK-immunoglobulin fusion proteins of the invention can be used as immunogens to produce anti-SphK antibodies in a subject, to purify SphK binding molecule, and in screening assays to identify molecules that inhibit the interaction of SphK with sphingosine or an SphK binding molecule.

An SphK chimeric or fusion protein of the invention can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, *e.g.*, by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including

WO 01/60990

PCT/US01/04789

automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers that give rise to complementary overhangs between two consecutive gene fragments that can subsequently be annealed and reamplified to generate a chimeric gene sequence (*see, e.g.*, Ausubel, *et al.* (eds.) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). An SphK-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the SphK protein.

SphK Agonists and Antagonists

The invention also pertains to variants of the SphK proteins that function as either SphK agonists (*i.e.*, mimetics) or as SphK antagonists. Variants of the SphK protein can be generated by mutagenesis (*e.g.*, discrete point mutation or truncation of the SphK protein). An agonist of the SphK protein can retain substantially the same, or a subset of, the biological activities of the naturally occurring form of the SphK protein. An antagonist of the SphK protein can inhibit one or more of the activities of the naturally occurring form of the SphK protein by, for example, competitively binding to a downstream or upstream member of a cellular signaling cascade which includes the SphK protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the SphK proteins.

Variants of the SphK proteins that function as either SphK agonists (*i.e.*, mimetics) or as SphK antagonists can be identified by screening combinatorial libraries of mutants (*e.g.*, truncation mutants) of the SphK proteins for SphK protein agonist or antagonist activity. In one embodiment, a variegated library of SphK variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of SphK variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential SphK sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display) containing the set of SphK sequences therein. There are a variety of methods that can be used to produce libraries of potential SphK variants from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential SphK sequences. Methods for

WO 01/60990

PCT/US01/04789

F_{ab} expression library. In general, antibody molecules obtained from humans relates to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another by the nature of the heavy chain present in the molecule. Certain classes have subclasses as well, such as IgG₁, IgG₂, and others. Furthermore, in humans, the light chain may be a kappa chain or a lambda chain. Reference herein to antibodies includes a reference to all such classes, subclasses and types of human antibody species.

An isolated protein of the invention intended to serve as an antigen, or a portion or fragment thereof, can be used as an immunogen to generate antibodies that immunospecifically bind the antigen, using standard techniques for polyclonal and monoclonal antibody preparation. The full-length protein can be used or, alternatively, the invention provides antigenic peptide fragments of the antigen for use as immunogens. An antigenic peptide fragment comprises at least 6 amino acid residues of the amino acid sequence of the full length protein, such as an amino acid sequence shown in SEQ ID NO:2, 4, 6, 8, 10, 12, 13, 14, or 15 and encompasses an epitope thereof such that an antibody raised against the peptide forms a specific immune complex with the full length protein or with any fragment that contains the epitope. Preferably, the antigenic peptide comprises at least 10 amino acid residues, or at least 15 amino acid residues, or at least 20 amino acid residues, or at least 30 amino acid residues. Preferred epitopes encompassed by the antigenic peptide are regions of the protein that are located on its surface; commonly these are hydrophilic regions.

In certain embodiments of the invention, at least one epitope encompassed by the antigenic peptide is a region of SphK that is located on the surface of the protein, e.g., a hydrophilic region. A hydrophobicity analysis of the SphK protein sequence will indicate which regions of a SphK polypeptide are particularly hydrophilic and, therefore, are likely to encode surface residues useful for targeting antibody production. As a means for targeting antibody production, hydropathy plots showing regions of hydrophilicity and hydrophobicity may be generated by any method well known in the art, including, for example, the Kyte Doolittle or the Hopp Woods methods, either with or without Fourier transformation. See, e.g., Hopp and Woods, 1981, *Proc. Nat. Acad. Sci. USA* 78: 3824-3828; Kyte and Doolittle 1982, *J. Mol. Biol.* 157: 105-142, each incorporated herein by reference in their entirety. Antibodies that are specific for one or more domains within an antigenic protein, or derivatives, fragments, analogs or homologs thereof, are also provided herein.

A protein of the invention, or a derivative, fragment, analog, homolog or ortholog thereof, may be utilized as an immunogen in the generation of antibodies that immunospecifically bind these protein components.

WO 01/60990

PCT/US01/04789

Various procedures known within the art may be used for the production of polyclonal or monoclonal antibodies directed against a protein of the invention, or against derivatives, fragments, analogs homologs or orthologs thereof (see, for example, Antibodies: A Laboratory Manual, Harlow E, and Lane D, 1988, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, incorporated herein by reference). Some of these antibodies are discussed below.

1. Polyclonal Antibodies

For the production of polyclonal antibodies, various suitable host animals (e.g., rabbit, goat, mouse or other mammal) may be immunized by one or more injections with the native protein, a synthetic variant thereof, or a derivative of the foregoing. An appropriate immunogenic preparation can contain, for example, the naturally occurring immunogenic protein, a chemically synthesized polypeptide representing the immunogenic protein, or a recombinantly expressed immunogenic protein. Furthermore, the protein may be conjugated to a second protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. The preparation can further include an adjuvant. Various adjuvants used to increase the immunological response include, but are not limited to, Freund's (complete and incomplete), mineral gels (e.g., aluminum hydroxide), surface active substances (e.g., lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, etc.), adjuvants usable in humans such as Bacille Calmette-Guerin and Corynebacterium parvum, or similar immunostimulatory agents. Additional examples of adjuvants which can be employed include MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate).

The polyclonal antibody molecules directed against the immunogenic protein can be isolated from the mammal (e.g., from the blood) and further purified by well known techniques, such as affinity chromatography using protein A or protein G, which provide primarily the IgG fraction of immune serum. Subsequently, or alternatively, the specific antigen that is the target of the immunoglobulin sought, or an epitope thereof, may be immobilized on a column to purify the immune specific antibody by immunoaffinity chromatography. Purification of immunoglobulins is discussed, for example, by D. Wilkinson (The Scientist, published by The Scientist, Inc., Philadelphia PA, Vol. 14, No. 8 (April 17, 2000), pp. 25-28).

2. Monoclonal Antibodies

The term "monoclonal antibody" (MAb) or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one molecular species of

WO 01/60990

PCT/US01/04789

antibody molecule consisting of a unique light chain gene product and a unique heavy chain gene product. In particular, the complementarity determining regions (CDRs) of the monoclonal antibody are identical in all the molecules of the population. MAbs thus contain an antigen binding site capable of immunoreacting with a particular epitope of the antigen characterized by
5 a unique binding affinity for it.

Monoclonal antibodies can be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind
10 to the immunizing agent. Alternatively, the lymphocytes can be immunized in vitro.

The immunizing agent will typically include the protein antigen, a fragment thereof or a fusion protein thereof. Generally, either peripheral blood lymphocytes are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a
15 suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells can be cultured in a suitable culture medium that preferably contains one or
20 more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma
30 and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

The culture medium in which the hybridoma cells are cultured can then be assayed for
35 the presence of monoclonal antibodies directed against the antigen. Preferably, the binding

WO 01/60990

PCT/US01/04789

specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the
 5 Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980). It is an objective, especially important in therapeutic applications of monoclonal antibodies, to identify antibodies having a high degree of specificity and a high binding affinity for the target antigen.

After the desired hybridoma cells are identified, the clones can be subcloned by limiting dilution procedures and grown by standard methods (Goding, 1986). Suitable culture media for
 10 this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells can be grown *in vivo* as ascites in a mammal.

The monoclonal antibodies secreted by the subclones can be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis,
 15 dialysis, or affinity chromatography.

The monoclonal antibodies can also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and
 20 light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also can be modified, for
 25 example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, Nature 368, 812-13 (1994)) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can
 30 be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

3. Humanized Antibodies

The antibodies directed against the protein antigens of the invention can further comprise humanized antibodies or human antibodies. These antibodies are suitable for administration to
 35 humans without engendering an immune response by the human against the administered

WO 01/60990

PCT/US01/04789

immunoglobulin. Humanized forms of antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) that are principally comprised of the sequence of a human immunoglobulin, and contain minimal sequence derived from a non-human immunoglobulin.

5 Humanization can be performed following the method of Winter and co-workers (Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeven et al., *Science*, 239:1534-1536 (1988)), by substituting rodent CDRs or CDR' sequences for the corresponding sequences of a human antibody. (See also U.S. Patent No. 5,225,539.) In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding

10 non-human residues. Humanized antibodies can also comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework regions are those of a

15 human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin (Jones et al., 1986; Riechmann et al., 1988; and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992)).

4. Human Antibodies

20 Fully human antibodies essentially relate to antibody molecules in which the entire sequence of both the light chain and the heavy chain, including the CDRs, arise from human genes. Such antibodies are termed "human antibodies"; or "fully human antibodies" herein. Human monoclonal antibodies can be prepared by the trioma technique; the human B-cell hybridoma technique (see Kozbor, et al., 1983 *Immunol Today* 4: 72) and the EBV hybridoma

25 technique to produce human monoclonal antibodies (see Cole, et al., 1985 In: *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96). Human monoclonal antibodies may be utilized in the practice of the present invention and may be produced by using human hybridomas (see Cote, et al., 1983. *Proc Natl Acad Sci USA* 80: 2026-2030) or by transforming human B-cells with Epstein Barr Virus in vitro (see Cole, et al., 1985 In:

30 *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96).

In addition, human antibodies can also be produced using additional techniques, including phage display libraries (Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)). Similarly, human antibodies can be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the

WO 01/60990

PCT/US01/04789

endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in Marks et al. (Bio/Technology 10, 779-783 (1992)); Lonberg et al. (Nature 368 856-859 (1994)); Morrison (Nature 368, 812-13 (1994)); Fishwild et al. (Nature Biotechnology 14, 845-51 (1996)); Neuberger (Nature Biotechnology 14, 826 (1996)); and Lonberg and Huszar (Intern. Rev. Immunol. 13 65-93 (1995)).

Human antibodies may additionally be produced using transgenic nonhuman animals that are modified so as to produce fully human antibodies rather than the animal's endogenous antibodies in response to challenge by an antigen. (See PCT publication WO94/02602). The endogenous genes encoding the heavy and light immunoglobulin chains in the nonhuman host have been incapacitated, and active loci encoding human heavy and light chain immunoglobulins are inserted into the host's genome. The human genes are incorporated, for example, using yeast artificial chromosomes containing the requisite human DNA segments. An animal which provides all the desired modifications is then obtained as progeny by crossbreeding intermediate transgenic animals containing fewer than the full complement of the modifications. The preferred embodiment of such a nonhuman animal is a mouse, and is termed the XenomouseTM as disclosed in PCT publications WO 96/33735 and WO 96/34096. This animal produces B cells that secrete fully human immunoglobulins. The antibodies can be obtained directly from the animal after immunization with an immunogen of interest, as, for example, a preparation of a polyclonal antibody, or alternatively from immortalized B cells, derived from the animal, such as hybridomas producing monoclonal antibodies. Additionally, the genes encoding the immunoglobulins with human variable regions can be recovered and expressed to obtain the antibodies directly, or can be further modified to obtain analogs of antibodies such as, for example, single chain Fv molecules.

An example of a method of producing a nonhuman host, exemplified as a mouse, lacking expression of an endogenous immunoglobulin heavy chain is disclosed in U.S. Patent No. 5,939,598. It can be obtained by a method including deleting the J segment genes from at least one endogenous heavy chain locus in an embryonic stem cell to prevent rearrangement of the locus and to prevent formation of a transcript of a rearranged immunoglobulin heavy chain locus, the deletion being effected by a targeting vector containing a gene encoding a selectable marker; and producing from the embryonic stem cell a transgenic mouse whose somatic and germ cells contain the gene encoding the selectable marker.

WO 01/60990

PCT/US01/04789

A method for producing an antibody of interest, such as a human antibody, is disclosed in U.S. Patent No. 5,916,771. It includes introducing an expression vector that contains a nucleotide sequence encoding a heavy chain into one mammalian host cell in culture, introducing an expression vector containing a nucleotide sequence encoding a light chain into
 5 another mammalian host cell, and fusing the two cells to form a hybrid cell. The hybrid cell expresses an antibody containing the heavy chain and the light chain.

In a further improvement on this procedure, a method for identifying a clinically relevant epitope on an immunogen, and a correlative method for selecting an antibody that binds immunospecifically to the relevant epitope with high affinity, are disclosed in PCT publication
 10 WO 99/53049.

5. F_{ab} Fragments and Single Chain Antibodies

According to the invention, techniques can be adapted for the production of single-chain antibodies specific to an antigenic protein of the invention (see e.g., U.S. Patent No. 4,946,778). In addition, methods can be adapted for the construction of F_{ab} expression libraries (see e.g.,
 15 Huse, et al., 1989 Science 246: 1275-1281) to allow rapid and effective identification of monoclonal F_{ab} fragments with the desired specificity for a protein or derivatives, fragments, analogs or homologs thereof. Antibody fragments that contain the idiotypes to a protein antigen may be produced by techniques known in the art including, but not limited to: (i) an F_{(ab)₂} fragment produced by pepsin digestion of an antibody molecule; (ii) an F_{ab} fragment generated
 20 by reducing the disulfide bridges of an F_{(ab)₂} fragment; (iii) an F_{ab} fragment generated by the treatment of the antibody molecule with papain and a reducing agent and (iv) F_v fragments.

6. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the
 25 binding specificities is for an antigenic protein of the invention. The second binding target is any other antigen, and advantageously is a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different
 30 specificities (Milstein and Cuello, *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity

WO 01/60990

PCT/US01/04789

chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker et al., EMBO J., 10:3655-3659 (1991).

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology, 121:210 (1986).

According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers that are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., Science 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Additionally, Fab' fragments can be directly recovered from E. coli and chemically coupled to form bispecific antibodies. Shalaby et al., J. Exp. Med. 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment

WO 01/60990

PCT/US01/04789

No. 4,676,980), and for treatment of HIV infection (WO 91/00360; WO 92/200373; EP 03089).

It is contemplated that the antibodies can be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins can be constructed using a disulfide exchange reaction or by forming a thioether bond.

- 5 Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

8. Effector Function Engineering

It can be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) can be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated can have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., *J. Exp Med.*, **176**: 1191-1195 (1992) and Shopes, *J. Immunol.*, **148**: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity can also be prepared using heterobifunctional cross-linkers as described in Wolff et al. *Cancer Research*, **53**: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and can thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., *Anti-Cancer Drug Design*, **3**: 219-230 (1989).

9. Immunoconjugates

20 The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolacca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, croton, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL),

WO 01/60990

PCT/US01/04789

active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a
 5 ricin immunotoxin can be prepared as described in Vitetta et al., Science, **238**: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

In another embodiment, the antibody can be conjugated to a "receptor" (such
 10 streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is in turn conjugated to a cytotoxic agent.

10. Immunoliposomes

15 The antibodies disclosed herein can also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., Proc. Natl. Acad. Sci. USA, **82**: 3688 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA, **77**: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

20 Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin et al., J. Biol. Chem., **257**:
 25 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See Gabizon et al., J. National Cancer Inst., **81**(19): 1484 (1989).

11. Diagnostic Applications of Antibodies Directed Against the Proteins of the Invention

Antibodies directed against a protein of the invention may be used in methods known
 30 within the art relating to the localization and/or quantitation of the protein (e.g., for use in measuring levels of the protein within appropriate physiological samples, for use in diagnostic methods, for use in imaging the protein, and the like). In a given embodiment, antibodies against the proteins, or derivatives, fragments, analogs or homologs thereof, that contain the antigen binding domain, are utilized as pharmacologically-active compounds (see below).

WO 01/60990

PCT/US01/04789

An antibody specific for a protein of the invention can be used to isolate the protein by standard techniques, such as immunoaffinity chromatography or immunoprecipitation. Such an antibody can facilitate the purification of the natural protein antigen from cells and of recombinantly produced antigen expressed in host cells. Moreover, such an antibody can be used to detect the antigenic protein (e.g., in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the antigenic protein. Antibodies directed against the protein can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, e.g., to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

12. Antibody Therapeutics

Antibodies of the invention, including polyclonal, monoclonal, humanized and fully human antibodies, may be used as therapeutic agents. Such agents will generally be employed to treat or prevent a disease or pathology in a subject. An antibody preparation, preferably one having high specificity and high affinity for its target antigen, is administered to the subject and will generally have an effect due to its binding with the target. Such an effect may be one of two kinds, depending on the specific nature of the interaction between the given antibody molecule and the target antigen in question. In the first instance, administration of the antibody may abrogate or inhibit the binding of the target with an endogenous ligand to which it naturally binds. In this case, the antibody binds to the target and masks a binding site of the naturally occurring ligand, wherein the ligand serves as an effector molecule. Thus the receptor mediates a signal transduction pathway for which ligand is responsible.

Alternatively, the effect may be one in which the antibody elicits a physiological result by virtue of binding to an effector binding site on the target molecule. In this case the target, a receptor having an endogenous ligand that may be absent or defective in the disease or pathology, binds the antibody as a surrogate effector ligand, initiating a receptor-based signal transduction event by the receptor.

WO 01/60990

PCT/US01/04789

A therapeutically effective amount of an antibody of the invention relates generally to the amount needed to achieve a therapeutic objective. As noted above, this may be a binding interaction between the antibody and its target antigen that, in certain cases, interferes with the functioning of the target, and in other cases, promotes a physiological response. The amount
5 required to be administered will furthermore depend on the binding affinity of the antibody for its specific antigen, and will also depend on the rate at which an administered antibody is depleted from the free volume other subject to which it is administered. Common ranges for therapeutically effective dosing of an antibody or antibody fragment of the invention may be, by way of nonlimiting example, from about 0.1 mg/kg body weight to about 50 mg/kg body weight.
10 Common dosing frequencies may range, for example, from twice daily to once a week.

13. Pharmaceutical Compositions of Antibodies

Antibodies specifically binding a protein of the invention, as well as other molecules identified by the screening assays disclosed herein, can be administered for the treatment of various disorders in the form of pharmaceutical compositions. Principles and considerations
15 involved in preparing such compositions, as well as guidance in the choice of components are provided, for example, in Remington : The Science And Practice Of Pharmacy 19th ed. (Alfonso R. Gennaro, et al., editors) Mack Pub. Co., Easton, Pa. : 1995; Drug Absorption Enhancement : Concepts, Possibilities, Limitations, And Trends, Harwood Academic Publishers, Langhorne, Pa., 1994; and Peptide And Protein Drug Delivery (Advances In Parenteral Sciences, Vol. 4),
20 1991, M. Dekker, New York.

If the antigenic protein is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is
25 preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, e.g., Marasco et al., Proc. Natl. Acad. Sci. USA, 90: 7889-7893 (1993). The formulation herein can also contain more than one active compound as necessary for the particular indication being
30 treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition can comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

WO 01/60990

PCT/US01/04789

The active ingredients can also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations can be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

14. ELISA Assay

An agent for detecting a SphK protein is an antibody capable of binding to the protein, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g., F_{ab} or $F_{(ab)2}$) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently-labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently-labeled streptavidin. The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. Included within the usage of the term "biological sample", therefore, is blood and a fraction or component of blood including blood serum, blood plasma, or lymph. That is, the detection method of the invention can be used to detect an SphK mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of an analyte mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of SphK protein

WO 01/60990

PCT/US01/04789

include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations, and immunofluorescence. *In vitro* techniques for detection of SphK genomic DNA include Southern hybridizations. Procedures for conducting immunoassays are described, for example in "ELISA: Theory and Practice: Methods in Molecular Biology", Vol. 42, J. R. Crowther (Ed.) Human Press, Totowa, NJ, 1995; "Immunoassay", E. Diamandis and T. Christopoulos, Academic Press, Inc., San Diego, CA, 1996; and "Practice and Theory of Enzyme Immunoassays", P. Tijssen, Elsevier Science Publishers, Amsterdam, 1985. Furthermore, *in vivo* techniques for detection of SphK protein include introducing into a subject a labeled anti-SphK protein antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

SphK Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding an SphK protein, or derivatives, fragments, analogs or homologs thereof. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively-linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, that is operatively-linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably-linked" is

WO 01/60990

PCT/US01/04789

intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner that allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell).

The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Regulatory sequences include those that direct constitutive expression of a nucleotide sequence in many types of host cell and those that direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention may be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (*e.g.*, SphK proteins, mutant forms of SphK proteins, fusion proteins, etc.).

The recombinant expression vectors of the invention may be designed for expression of SphK proteins in prokaryotic or eukaryotic cells. For example, SphK proteins may be expressed in bacterial cells such as *Escherichia coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *Escherichia coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: (i) to increase expression of recombinant protein; (ii) to increase the solubility of the recombinant protein; and (iii) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith and Johnson, 1988. *Gene* 67: 31-40), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.) that fuse glutathione

WO 01/60990

PCT/US01/04789

S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amrann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, GENE EXPRESSION
5 TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990) 60-89).

One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein. See, e.g., Gottesman, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY
10 185, Academic Press, San Diego, Calif. (1990) 119-128. Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (see, e.g., Wada, *et al.*, 1992. *Nucl. Acids Res.* 20: 2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

15 In another embodiment, the SphK expression vector is a yeast expression vector. Examples of vectors for expression in yeast *Saccharomyces cerevisiae* include pYepSec1 (Baldari, *et al.*, 1987. *EMBO J.* 6: 229-234), pMFa (Kurjan and Herskowitz, 1982. *Cell* 30: 933-943), pJRY88 (Schultz *et al.*, 1987. *Gene* 54: 113-123), pYES2 (Invitrogen Corporation, San Diego, Calif.), and picZ (InVitrogen Corp, San Diego, Calif.).

20 Alternatively, SphK can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., SF9 cells) include the pAc series (Smith, *et al.*, 1983. *Mol. Cell. Biol.* 3: 2156-2165) and the pVL series (Lucklow and Summers, 1989. *Virology* 170: 31-39).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian
25 cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, 1987. *Nature* 329: 840) and pMT2PC (Kaufman, *et al.*, 1987. *EMBO J.* 6: 187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, adenovirus 2, cytomegalovirus, and simian virus 40. For other suitable
30 expression systems for both prokaryotic and eukaryotic cells see, e.g., Chapters 16 and 17 of Sambrook, *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory
35

WO 01/60990

PCT/US01/04789

elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert, *et al.*, 1987. *Genes Dev.* 1: 268-277), lymphoid-specific promoters (Calame and Eaton, 1988. *Adv. Immunol.* 43: 235-275), in particular promoters of T cell receptors (Winoto and Baltimore, 1989. *EMBO J.* 8: 729-733) and immunoglobulins (Banerji, *et al.*, 1983. *Cell* 33: 729-740; Queen and Baltimore, 1983. *Cell* 33: 741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle, 1989. *Proc. Natl. Acad. Sci. USA* 86: 5473-5477), pancreas-specific promoters (Edlund, *et al.*, 1985. *Science* 230: 912-916), and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S. Pat. No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, *e.g.*, the murine hox promoters (Kessel and Gruss, 1990. *Science* 249: 374-379) and the α -fetoprotein promoter (Campes and Tilghman, 1989. *Genes Dev.* 3: 537-546).

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively-linked to a regulatory sequence in a manner that allows for expression (by transcription of the DNA molecule) of an RNA molecule that is antisense to SphK mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen that direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen that direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes *see, e.g.*, Weintraub, *et al.*, "Antisense RNA as a molecular tool for genetic analysis," *Reviews-Trends in Genetics*, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but also to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, SphK protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as

WO 01/60990

PCT/US01/04789

Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and
 5 "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (e.g., DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (MOLECULAR CLONING: A LABORATORY MANUAL, 2nd ed., Cold Spring Harbor Laboratory,
 10 Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that
 15 encodes a selectable marker (e.g., resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Various selectable markers include those that confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be introduced into a host cell on the same vector as that encoding SphK or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid
 20 can be identified by drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (*i.e.*, express) SphK protein. Accordingly, the invention further provides methods for producing SphK protein using the host cells of the invention. In one embodiment,
 25 the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding SphK protein has been introduced) in a suitable medium such that SphK protein is produced. In another embodiment, the method further comprises isolating SphK protein from the medium or the host cell.

Transgenic SphK Animals

30 The host cells of the invention can also be used to produce non-human transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which SphK protein-coding sequences have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous SphK sequences have been introduced into their genome or homologous recombinant animals in which

WO 01/60990

PCT/US01/04789

(e.g., the cDNA of SEQ ID NO:1 or 5), but more preferably, is a non-human homologue of a human SphK gene. For example, a mouse homologue of human SphK gene can be used to construct a homologous recombination vector suitable for altering an endogenous SphK gene in the mouse genome. In one embodiment, the vector is designed such that, upon homologous recombination, the endogenous SphK gene is functionally disrupted (i.e., no longer encodes a functional protein; also referred to as a "knock out" vector).

Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous SphK gene is mutated or otherwise altered but still encodes functional protein (e.g., the upstream regulatory region can be altered to thereby alter the expression of the endogenous SphK protein). In the homologous recombination vector, the altered portion of the SphK gene is flanked at its 5'- and 3'-termini by additional nucleic acid of the SphK gene to allow for homologous recombination to occur between the exogenous SphK gene carried by the vector and an endogenous SphK gene in an embryonic stem cell. The additional flanking SphK nucleic acid is of sufficient length for successful homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5'- and 3'-termini) are included in the vector. See, e.g., Thomas, *et al.*, 1987. *Cell* 51: 503 for a description of homologous recombination vectors. The vector is then introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced SphK gene has homologously-recombined with the endogenous SphK gene are selected. See, e.g., Li, *et al.*, 1992. *Cell* 69: 915.

The selected cells are then injected into a blastocyst of an animal (e.g., a mouse) to form aggregation chimeras. See, e.g., Bradley, 1987. In: *TERATOCARCINOMAS AND EMBRYONIC STEM CELLS: A PRACTICAL APPROACH*, Robertson, ed. IRL, Oxford, pp. 113-152. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harboring the homologously-recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously-recombined DNA by germline transmission of the transgene. Methods for constructing homologous recombination vectors and homologous recombinant animals are described further in Bradley, 1991. *Curr. Opin. Biotechnol.* 2: 823-829; PCT International Publication Nos.: WO 90/11354; WO 91/01140; WO 92/0968; and WO 93/04169.

In another embodiment, transgenic non-humans animals can be produced that contain selected systems that allow for regulated expression of the transgene. One example of such a system is the cre/loxP recombinase system of bacteriophage P1. For a description of the cre/loxP recombinase system, See, e.g., Lakso, *et al.*, 1992. *Proc. Natl. Acad. Sci. USA* 89: 6232-6236. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae*. See, O'Gorman, *et al.*, 1991. *Science* 251:1351-1355. If a cre/loxP

WO 01/60990

PCT/US01/04789

recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other
5 containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, *et al.*, 1997. *Nature* 385: 810-813. In brief, a cell (e.g., a somatic cell) from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can then be fused, e.g., through the use of electrical
10 pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyte and then transferred to pseudopregnant female foster animal. The offspring borne of this female foster animal will be a clone of the animal from which the cell (e.g., the somatic cell) is isolated.

15 **Pharmaceutical Compositions**

The SphK nucleic acid molecules, SphK proteins, and anti-SphK antibodies (also referred to herein as "active compounds") of the invention, and derivatives, fragments, analogs and homologs thereof, can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or
20 antibody and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Suitable carriers are described in the most recent edition of Remington's Pharmaceutical Sciences, a standard reference text in the field,
25 which is incorporated herein by reference. Preferred examples of such carriers or diluents include, but are not limited to, water, saline, finger's solutions, dextrose solution, and 5% human serum albumin. Liposomes and non-aqueous vehicles such as fixed oils may also be used. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound,
30 use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (i.e., topical),

WO 01/60990

PCT/US01/04789

transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (e.g., an SphK protein or anti-SphK antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of

WO 01/60990

PCT/US01/04789

the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser that contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including

WO 01/60990

PCT/US01/04789

liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (*see, e.g.*, U.S. Patent No. 5,328,470) or by stereotactic injection (*see, e.g.*, Chen, *et al.*, 1994. *Proc. Natl. Acad. Sci. USA* 91: 3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, *e.g.*, retroviral vectors, the pharmaceutical preparation can include one or more cells that produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

Screening and Detection Methods

The isolated nucleic acid molecules of the invention can be used to express SphK protein (*e.g.*, via a recombinant expression vector in a host cell in gene therapy applications), to detect SphK mRNA (*e.g.*, in a biological sample) or a genetic lesion in an SphK gene, and to modulate SphK activity, as described further, below. In addition, the SphK proteins can be used to screen drugs or compounds that modulate the SphK protein activity or expression as well as to treat disorders characterized by insufficient or excessive production of SphK protein or production of SphK protein forms that have decreased or aberrant activity compared to SphK wild-type protein. In addition, the anti-SphK antibodies of the invention can be used to detect and isolate SphK proteins and modulate SphK activity.

WO 01/60990

PCT/US01/04789

The invention further pertains to novel agents identified by the screening assays described herein and uses thereof for treatments as described, *supra*.

Screening Assays

The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*, peptides, peptidomimetics, small molecules or other drugs) that bind to SphK proteins or have a stimulatory or inhibitory effect on, *e.g.*, SphK protein expression or SphK protein activity. The invention also includes compounds identified in the screening assays described herein.

In one embodiment, the invention provides assays for screening candidate or test compounds that bind to or modulate the activity of SphK protein or polypeptide or biologically-active portion thereof. The test compounds of the invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds. *See, e.g.*, Lam, 1997.

Anticancer Drug Design 12: 145.

A "small molecule" as used herein, is meant to refer to a composition that has a molecular weight of less than about 5 kD and most preferably less than about 4 kD. Small molecules can be, *e.g.*, nucleic acids, peptides, polypeptides, peptidomimetics, carbohydrates, lipids or other organic or inorganic molecules. Libraries of chemical and/or biological mixtures, such as fungal, bacterial, or algal extracts, are known in the art and can be screened with any of the assays of the invention.

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt, *et al.*, 1993. *Proc. Natl. Acad. Sci. U.S.A.* 90: 6909; Erb, *et al.*, 1994. *Proc. Natl. Acad. Sci. U.S.A.* 91: 11422; Zuckermann, *et al.*, 1994. *J. Med. Chem.* 37: 2678; Cho, *et al.*, 1993. *Science* 261: 1303; Carrell, *et al.*, 1994. *Angew. Chem. Int. Ed. Engl.* 33: 2059; Carrell, *et al.*, 1994. *Angew. Chem. Int. Ed. Engl.* 33: 2061; and Gallop, *et al.*, 1994. *J. Med. Chem.* 37: 1233.

Libraries of compounds may be presented in solution (*e.g.*, Houghten, 1992. *Biotechniques* 13: 412-421), or on beads (Lam, 1991. *Nature* 354: 82-84), on chips (Fodor, 1993. *Nature* 364: 555-556), bacteria (Ladner, U.S. Patent No. 5,223,409), spores (Ladner, U.S. Patent 5,233,409), plasmids (Cull, *et al.*, 1992. *Proc. Natl. Acad. Sci. USA* 89: 1865-1869) or on phage (Scott and Smith, 1990. *Science* 249: 386-390; Devlin, 1990. *Science* 249: 404-406;

WO 01/60990

PCT/US01/04789

Cwirla, *et al.*, 1990. *Proc. Natl. Acad. Sci. U.S.A.* 87: 6378-6382; Felici, 1991. *J. Mol. Biol.* 222: 301-310; Ladner, U.S. Patent No. 5,233,409.).

In one embodiment, an assay is a cell-based assay in which a cell that has been engineered to express a membrane-bound form of an SphK protein, or a biologically-active portion thereof, on the cell surface is contacted with a test compound and the ability of the test compound to bind to an SphK protein determined. The cell, for example, can be of mammalian origin or a yeast cell. Determining the ability of the test compound to bind to the SphK protein can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the SphK protein or biologically-active portion thereof can be determined by detecting the labeled compound in a complex. For example, test compounds can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, test compounds can be enzymatically-labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product. In one embodiment, the assay comprises contacting a cell which expresses a membrane-bound form of SphK protein, or a biologically-active portion thereof, on the cell surface with a known compound which binds SphK to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with an SphK protein, wherein determining the ability of the test compound to interact with an SphK protein comprises determining the ability of the test compound to preferentially bind to SphK protein or a biologically-active portion thereof as compared to the known compound.

In another embodiment, an assay is a cell-based assay comprising contacting a cell engineered to express a membrane-bound form of SphK protein, or a biologically-active portion thereof, on the cell surface with a test compound and determining the ability of the test compound to modulate (*e.g.*, stimulate or inhibit) the activity of the SphK protein or biologically-active portion thereof. Determining the ability of the test compound to modulate the activity of SphK or a biologically-active portion thereof can be accomplished, for example, by determining the ability of the SphK protein to phosphorylate sphingosine. Alternatively, the ability of SphK to bind a target molecule may be determined. As used herein, a "target molecule" is a molecule with which an SphK protein binds or interacts in nature. An SphK target molecule can be a non-SphK molecule or an SphK protein or polypeptide of the invention. In one embodiment, an SphK target molecule is a component of a signal transduction pathway that facilitates transduction of an extracellular signal through the cell membrane and into the

WO 01/60990

PCT/US01/04789

cell. The target, for example, can be a second intercellular protein that has catalytic activity or a protein that facilitates the association of downstream signaling molecules with SphK.

Determining the ability of the SphK protein to bind to or interact with an SphK target molecule can be accomplished by one of the methods described above for determining direct
5 binding. In one embodiment, determining the ability of the SphK protein to bind to or interact with an SphK target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (*i.e.* intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of the target an appropriate substrate, detecting the
10 induction of a reporter gene (comprising an SphK -responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a cellular response, for example, cell survival, cellular differentiation, or cell proliferation.

In yet another embodiment, an assay of the invention is a cell-free assay comprising contacting an SphK protein or biologically-active portion thereof with a test compound and
15 determining the ability of the test compound to bind to the SphK protein or biologically-active portion thereof. Binding of the test compound to the SphK protein can be determined either directly or indirectly as described above. In one such embodiment, the assay comprises contacting the SphK protein or biologically-active portion thereof with a known compound which binds SphK to form an assay mixture, contacting the assay mixture with a test compound,
20 and determining the ability of the test compound to interact with an SphK protein, wherein determining the ability of the test compound to interact with an SphK protein comprises determining the ability of the test compound to preferentially bind to SphK or biologically-active portion thereof as compared to the known compound.

In still another embodiment, an assay is a cell-free assay comprising contacting SphK
25 protein or biologically-active portion thereof with a test compound and determining the ability of the test compound to modulate (*e.g.* stimulate or inhibit) the activity of the SphK protein or biologically-active portion thereof (*e.g.*, the ability to phosphorylate sphingosine). Determining the ability of the test compound to modulate the activity of SphK can be accomplished, for example, by determining the ability of the SphK protein to bind to an SphK target molecule by
30 one of the methods described above for determining direct binding. In an alternative embodiment, determining the ability of the test compound to modulate the activity of SphK protein can be accomplished by determining the ability of the SphK protein further modulate an SphK target molecule. For example, the catalytic/enzymatic activity of the target molecule on an appropriate substrate can be determined as described, *supra*.

WO 01/60990

PCT/US01/04789

In yet another embodiment, the cell-free assay comprises contacting the SphK protein or biologically-active portion thereof with a known compound which binds SphK protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with an SphK protein, wherein determining the ability of the test
5 compound to interact with an SphK protein comprises determining the ability of the SphK protein to preferentially bind to or modulate the activity of an SphK target molecule.

In more than one embodiment of the above assay methods of the invention, it may be desirable to immobilize either SphK protein or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate
10 automation of the assay. Binding of a test compound to SphK protein, or interaction of SphK protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion
15 protein can be provided that adds a domain that allows one or both of the proteins to be bound to a matrix. For example, GST-SphK fusion proteins or GST-target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtiter plates, that are then combined with the test compound or the test
20 compound and either the non-adsorbed target protein or SphK protein, and the mixture is incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined
either directly or indirectly, for example, as described, *supra*. Alternatively, the complexes can be dissociated from the matrix, and the level of SphK protein binding or activity determined using standard techniques.

25 Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either the SphK protein or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated SphK protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well-known within the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.), and
30 immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with SphK protein or target molecules, but which do not interfere with binding of the SphK protein to its target molecule, can be derivatized to the wells of the plate, and unbound target or SphK protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized
35 complexes, include immunodetection of complexes using antibodies reactive with the SphK

WO 01/60990

PCT/US01/04789

protein or target molecule, as well as enzyme-linked assays that rely on detecting an enzymatic activity associated with the SphK protein or target molecule.

In another embodiment, modulators of SphK protein expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of SphK mRNA or protein in the cell is determined. The level of expression of SphK mRNA or protein in the presence of the candidate compound is compared to the level of expression of SphK mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of SphK mRNA or protein expression based upon this comparison. For example, when expression of SphK mRNA or protein is greater (*i.e.*, statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of SphK mRNA or protein expression. Alternatively, when expression of SphK mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of SphK mRNA or protein expression. The level of SphK mRNA or protein expression in the cells can be determined by methods described herein for detecting SphK mRNA or protein.

In yet another aspect of the invention, the SphK proteins can be used as "bait proteins" in a two-hybrid assay or three hybrid assay (*see, e.g.*, U.S. Patent No. 5,283,317; Zervos, *et al.*, 1993. *Cell* 72: 223-232; Madura, *et al.*, 1993. *J. Biol. Chem.* 268: 12046-12054; Bartel, *et al.*, 1993. *Biotechniques* 14: 920-924; Iwabuchi, *et al.*, 1993. *Oncogene* 8: 1693-1696; and Brent WO 94/10300), to identify other proteins that bind to or interact with SphK ("SphK-binding proteins" or "SphK-bp") and modulate SphK activity. Such SphK-binding proteins are also likely to be involved in the propagation of signals by the SphK proteins as, for example, upstream or downstream elements of the SphK pathway.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for SphK is fused to a gene encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming an SphK-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ) that is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the

WO 01/60990

PCT/US01/04789

functional transcription factor can be isolated and used to obtain the cloned gene that encodes the protein that interacts with SphK.

The invention further pertains to novel agents identified by the aforementioned screening assays and uses thereof for treatments as described herein.

5 *Detection Assays*

Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. By way of example, and not of limitation, these sequences can be used to: (i) map their respective genes on a chromosome; and, thus, locate gene regions associated with genetic disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. Some of these applications are described in the subsections, below.

Chromosome Mapping

Once the sequence (or a portion of the sequence) of a gene has been isolated, this sequence can be used to map the location of the gene on a chromosome. This process is called chromosome mapping. Accordingly, portions or fragments of the SphK sequences, SEQ ID NO:1, 3, 5, 7, or 9, can be used to map the location of the SphK genes, respectively, on a chromosome. The mapping of the SphK sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

Briefly, SphK genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in length) from the SphK sequences. Computer analysis of the SphK, sequences can be used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers can then be used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the SphK sequences will yield an amplified fragment.

Somatic cell hybrids are prepared by fusing somatic cells from different mammals (*e.g.*, human and mouse cells). As hybrids of human and mouse cells grow and divide, they gradually lose human chromosomes in random order, but retain the mouse chromosomes. By using media in which mouse cells cannot grow, because they lack a particular enzyme, but in which human cells can, the one human chromosome that contains the gene encoding the needed enzyme will be retained. By using various media, panels of hybrid cell lines can be established. Each cell line in a panel contains either a single human chromosome or a small number of human chromosomes, and a full set of mouse chromosomes, allowing easy mapping of individual genes to specific human chromosomes. *See, e.g., D'Eustachio, et al., 1983. Science 220: 919-924.*

WO 01/60990

PCT/US01/04789

Somatic cell hybrids containing only fragments of human chromosomes can also be produced by using human chromosomes with translocations and deletions.

5 PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular sequence to a particular chromosome. Three or more sequences can be assigned per day using a single thermal cycler. Using the SphK sequences to design oligonucleotide primers, sub-localization can be achieved with panels of fragments from specific chromosomes.

10 Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase chromosomal spread can further be used to provide a precise chromosomal location in one step. Chromosome spreads can be made using cells whose division has been blocked in metaphase by a chemical like colcemid that disrupts the mitotic spindle. The chromosomes can be treated briefly with trypsin, and then stained with Giemsa. A pattern of light and dark bands develops on each chromosome, so that the chromosomes can be identified individually. The FISH technique can be used with a DNA sequence as short as 500 or 600 bases. However, clones larger than 1,000 bases have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. Preferably 1,000 bases, and more preferably 2,000 bases, will suffice to get good results at a reasonable amount of time. For a review of this technique, see, Verma, *et al.*, HUMAN CHROMOSOMES: A MANUAL OF BASIC TECHNIQUES (Pergamon Press, New York 1988).

20 Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

25 Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, *e.g.*, in McKusick, MENDELIAN INHERITANCE IN MAN, available on-line through Johns Hopkins University Welch Medical Library). The relationship between genes and disease, mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, *e.g.*, Egeland, *et al.*, 1987. *Nature*, 30 325: 783-787.

35 Moreover, differences in the DNA sequences between individuals affected and unaffected with a disease associated with the SphK gene, can be determined. If a mutation is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease. Comparison of affected and

WO 01/60990

PCT/US01/04789

unaffected individuals generally involves first looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of genes from several individuals can be performed to confirm the presence of a mutation and to
5 distinguish mutations from polymorphisms.

Tissue Typing

The SphK sequences of the invention can also be used to identify individuals from minute biological samples. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for
10 identification. The sequences of the invention are useful as additional DNA markers for RFLP ("restriction fragment length polymorphisms," described in U.S. Patent No. 5,272,057).

Furthermore, the sequences of the invention can be used to provide an alternative technique that determines the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the SphK sequences described herein can be used to prepare two
15 PCR primers from the 5'- and 3'-termini of the sequences. These primers can then be used to amplify an individual's DNA and subsequently sequence it.

Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a unique set of such DNA sequences due to allelic differences. The sequences of the invention can be used to obtain such
20 identification sequences from individuals and from tissue. The human SphK sequences of the invention uniquely represent portions of the human genome. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. It is estimated that allelic variation between individual humans occurs with a frequency of about once per each 500 bases. Much of the allelic variation is due to single nucleotide
25 polymorphisms (SNPs), which include restriction fragment length polymorphisms (RFLPs).

Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the noncoding regions, fewer sequences are necessary to differentiate individuals. The noncoding sequences can comfortably provide
30 positive individual identification with a panel of perhaps 10 to 1,000 primers that each yield a noncoding amplified sequence of 100 bases. If predicted coding sequences, such as those in SEQ ID NO:1, 3, 5, 7, or 9, are used, a more appropriate number of primers for positive individual identification would be 500-2,000.

WO/01/60990

PCT/US01/04789

conditions to SphK mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays of the invention are described herein.

An agent for detecting SphK protein is an antibody capable of binding to SphK protein, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g., Fab or F(ab')₂) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently-labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently-labeled streptavidin. The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention can be used to detect SphK mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of SphK mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of SphK protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations, and immunofluorescence. *In vitro* techniques for detection of SphK genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of SphK protein include introducing into a subject a labeled anti-SphK antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

In one embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject.

In another embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting SphK protein, mRNA, or genomic DNA, such that the presence of SphK protein, mRNA or genomic DNA is detected in the biological sample, and comparing the presence of SphK protein, mRNA or genomic DNA in the control sample with the presence of SphK protein, mRNA or genomic DNA in the test sample.

The invention also encompasses kits for detecting the presence of SphK in a biological sample. For example, the kit can comprise: a labeled compound or agent capable of detecting SphK protein or mRNA in a biological sample; means for determining the amount of SphK in the sample; and means for comparing the amount of SphK in the sample with a standard. The

WO 01/60990

PCT/US01/04789

compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect SphK protein or nucleic acid.

Prognostic Assays

The diagnostic methods described herein can furthermore be utilized to identify subjects
 5 having or at risk of developing a disease or disorder associated with aberrant SphK expression or activity. For example, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk of developing a disorder associated with SphK protein, nucleic acid expression or activity. Alternatively, the prognostic assays can be utilized to identify a subject having or at risk for developing a disease
 10 or disorder. Thus, the invention provides a method for identifying a disease or disorder associated with aberrant SphK expression or activity in which a test sample is obtained from a subject and SphK protein or nucleic acid (e.g., mRNA, genomic DNA) is detected, wherein the presence of SphK protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant SphK expression or activity. As used
 15 herein, a "test sample" refers to a biological sample obtained from a subject of interest. For example, a test sample can be a biological fluid (e.g., serum), cell sample, or tissue.

Furthermore, the prognostic assays described herein can be used to determine whether a subject can be administered an agent (e.g., an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder
 20 associated with aberrant SphK expression or activity. For example, such methods can be used to determine whether a subject can be effectively treated with an agent for a disorder. Thus, the invention provides methods for determining whether a subject can be effectively treated with an agent for a disorder associated with aberrant SphK expression or activity in which a test sample is obtained and SphK protein or nucleic acid is detected (e.g., wherein the presence of SphK
 25 protein or nucleic acid is diagnostic for a subject that can be administered the agent to treat a disorder associated with aberrant SphK expression or activity).

The methods of the invention can also be used to detect genetic lesions in an SphK gene, thereby determining if a subject with the lesioned gene is at risk for a disorder characterized by aberrant cell proliferation and/or differentiation. In various embodiments, the methods include
 30 detecting, in a sample of cells from the subject, the presence or absence of a genetic lesion characterized by at least one of an alteration affecting the integrity of a gene encoding an SphK protein, or the misexpression of the SphK gene. For example, such genetic lesions can be detected by ascertaining the existence of at least one of: (i) a deletion of one or more nucleotides from an SphK gene; (ii) an addition of one or more nucleotides to an SphK gene; (iii) a
 35 substitution of one or more nucleotides of an SphK gene, (iv) a chromosomal rearrangement of

WO 01/60990

PCT/US01/04789

5,493,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

In other embodiments, genetic mutations in SphK can be identified by hybridizing a sample and control nucleic acids, e.g., DNA or RNA, to high-density arrays containing hundreds or thousands of oligonucleotide probes. See, e.g., Cronin, *et al.*, 1996. *Human Mutation* 7: 244-255; Kozal, *et al.*, 1996. *Nat. Med.* 2: 753-759. For example, genetic mutations in SphK can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin, *et al.*, *supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the SphK gene and detect mutations by comparing the sequence of the sample SphK with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxim and Gilbert, 1977. *Proc. Natl. Acad. Sci. USA* 74: 560 or Sanger, 1977. *Proc. Natl. Acad. Sci. USA* 74: 5463. It is also contemplated that any of a variety of automated sequencing procedures can be utilized when performing the diagnostic assays (see, e.g., Naeve, *et al.*, 1995. *Biotechniques* 19: 448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen, *et al.*, 1996. *Adv. Chromatography* 36: 127-162; and Griffin, *et al.*, 1993. *Appl. Biochem. Biotechnol.* 38: 147-159).

Other methods for detecting mutations in the SphK gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes. See, e.g., Myers, *et al.*, 1985. *Science* 230: 1242. In general, the art technique of "mismatch cleavage" starts by providing heteroduplexes of formed by hybridizing (labeled) RNA or DNA containing the wild-type SphK sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent that cleaves single-stranded regions of the duplex such as which will exist due to basepair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA hybrids treated with S₁ nuclease to enzymatically digest the mismatched regions. In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to digest

WO 01/60990

PCT/US01/04789

mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. *See, e.g., Cotton, et al., 1988. Proc. Natl. Acad. Sci. USA 85: 4397; Saleeba, et al., 1992. Methods Enzymol. 217: 286-295.* In an embodiment, the control DNA or RNA can be labeled for
5 detection.

In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in SphK cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves
10 A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches. *See, e.g., Hsu, et al., 1994. Carcinogenesis 15: 1657-1662.* According to an exemplary embodiment, a probe based on an SphK sequence, *e.g., a wild-type SphK sequence,* is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme, and the cleavage products, if any, can be detected from
15 electrophoresis protocols or the like. *See, e.g., U.S. Patent No. 5,459,039.*

In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in SphK genes. For example, single strand conformation polymorphism (SSCP) may be used to detect differences in electrophoretic mobility between mutant and wild type nucleic acids. *See, e.g., Orita, et al., 1989. Proc. Natl. Acad. Sci. USA: 86: 2766; Cotton, 1993. Mutat.*
20 *Res. 285: 125-144; Hayashi, 1992. Genet. Anal. Tech. Appl. 9: 73-79.* Single-stranded DNA fragments of sample and control SphK nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of
25 the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In one embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility. *See, e.g., Keen, et al., 1991. Trends Genet. 7: 5.*

In yet another embodiment, the movement of mutant or wild-type fragments in
30 polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (DGGE). *See, e.g., Myers, et al., 1985. Nature 313: 495.* When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient

WO 01/60990

PCT/US01/04789

to identify differences in the mobility of control and sample DNA. *See, e.g., Rosenbaum and Reissner, 1987. Biophys. Chem. 265: 12753.*

Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension. For example, oligonucleotide primers may be prepared in which the known mutation is placed centrally and then hybridized to target DNA under conditions that permit hybridization only if a perfect match is found. *See, e.g., Saiki, et al., 1986. Nature 324: 163; Saiki, et al., 1989. Proc. Natl. Acad. Sci. USA 86: 6230.* Such allele specific oligonucleotides are hybridized to PCR amplified target DNA or a number of different mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

Alternatively, allele specific amplification technology that depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization; *see, e.g., Gibbs, et al., 1989. Nucl. Acids Res. 17: 2437-2448*) or at the extreme 3'-terminus of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (*see, e.g., Prossner, 1993. Tibtech. 11: 238*). In addition it may be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection. *See, e.g., Gasparini, et al., 1992. Mol. Cell Probes 6: 1.* It is anticipated that in certain embodiments amplification may also be performed using *Taq* ligase for amplification. *See, e.g., Barany, 1991. Proc. Natl. Acad. Sci. USA 88: 189.* In such cases, ligation will occur only if there is a perfect match at the 3'-terminus of the 5' sequence, making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, *e.g.,* in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving an SphK gene.

Furthermore, any cell type or tissue in which SphK is expressed may be utilized in the prognostic assays described herein.

30 *Pharmacogenomics*

Agents, or modulators that have a stimulatory or inhibitory effect on SphK activity (*e.g.,* SphK gene expression), as identified by a screening assay described herein can be administered to individuals to treat (prophylactically or therapeutically) disorders (The disorders include cell proliferative disorders such as cancers, ischemia, or restenosis. In conjunction with such treatment, the pharmacogenomics (*i.e.,* the study of the relationship between an individual's

WO 01/60990

PCT/US01/04789

genotype and that individual's response to a foreign compound or drug) of the individual may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, the pharmacogenomics of the individual permits the selection of effective agents (e.g., drugs) for prophylactic or therapeutic treatments based on a consideration of the individual's genotype. Such pharmacogenomics can further be used to determine appropriate dosages and therapeutic regimens. Accordingly, the activity of SphK protein, expression of SphK nucleic acid, or mutation content of SphK genes in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See e.g., Eichelbaum, 1996. *Clin. Exp. Pharmacol. Physiol.*, 23: 983-985; Linder, 1997. *Clin. Chem.*, 43: 254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare defects or as polymorphisms. For example, glucose-6-phosphate dehydrogenase (G6PD) deficiency is a common inherited enzymopathy in which the main clinical complication is hemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic polymorphisms of drug metabolizing enzymes (e.g., N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses. If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed metabolite morphine. At the other extreme are the so called ultra-rapid

WO 01/60990

PCT/US01/04789

metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Thus, the activity of SphK protein, expression of SphK nucleic acid, or mutation content of SphK genes in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual. In addition, pharmacogenetic studies can be used to apply genotyping of polymorphic alleles encoding drug-metabolizing enzymes to the identification of an individual's drug responsiveness phenotype. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with an SphK modulator, such as a modulator identified by one of the exemplary screening assays described herein.

Monitoring of Effects During Clinical Trials

Monitoring the influence of agents (e.g., drugs, compounds) on the expression or activity of SphK (e.g., the ability to modulate aberrant cell proliferation and/or differentiation) can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a screening assay as described herein to increase SphK gene expression, protein levels, or upregulate SphK activity, can be monitored in clinical trails of subjects exhibiting decreased SphK gene expression, protein levels, or downregulated SphK activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease SphK gene expression, protein levels, or downregulate SphK activity, can be monitored in clinical trails of subjects exhibiting increased SphK gene expression, protein levels, or upregulated SphK activity. In such clinical trials, the expression or activity of SphK and, preferably, other genes that have been implicated in, for example, a cellular proliferation or immune disorder can be used as a "read out" or markers of the immune responsiveness of a particular cell.

By way of example, and not of limitation, genes, including SphK, that are modulated in cells by treatment with an agent (e.g., compound, drug or small molecule) that modulates SphK activity (e.g., identified in a screening assay as described herein) can be identified. Thus, to study the effect of agents on cellular proliferation disorders, for example, in a clinical trial, cells can be isolated and RNA prepared and analyzed for the levels of expression of SphK and other genes implicated in the disorder. The levels of gene expression (i.e., a gene expression pattern) can be quantified by Northern blot analysis or RT-PCR, as described herein, or alternatively by measuring the amount of protein produced, by one of the methods as described herein, or by measuring the levels of activity of SphK or other genes. In this manner, the gene expression pattern can serve as a marker, indicative of the physiological response of the cells to the agent.

WO 01/60990

PCT/US01/04789

Accordingly, this response state may be determined before, and at various points during, treatment of the individual with the agent.

In one embodiment, the invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (e.g., an agonist, antagonist, protein, peptide, peptidomimetic, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of an SphK protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the SphK protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the SphK protein, mRNA, or genomic DNA in the pre-administration sample with the SphK protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent may be desirable to increase the expression or activity of SphK to higher levels than detected, *i.e.*, to increase the effectiveness of the agent. Alternatively, decreased administration of the agent may be desirable to decrease expression or activity of SphK to lower levels than detected, *i.e.*, to decrease the effectiveness of the agent.

Methods of Treatment

The invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant SphK expression or activity. The disorders include cell proliferative disorders such as cancer, ischemia, and restenosis. These methods of treatment will be discussed more fully, below.

Disease and Disorders

Diseases and disorders that are characterized by increased (relative to a subject not suffering from the disease or disorder) levels or biological activity may be treated with therapeutics that antagonize (*i.e.*, reduce or inhibit) activity. This decrease in activity may come about in a variety of ways, for example: (1) by decreasing the copies of the gene in the cell (amplifier); (2) by decreasing transcription of the SphK gene (transcription down-regulators); (3) by decreasing the translation of SphK mRNA into protein (translation down-regulators); or (4) by decreasing the activity of SphK itself (antagonists).

Expression of SphK inhibits apoptosis. Xia *et al.*, *J Biol Chem* 1999 Nov 26;274(48):34499-505. Cancer cells expressing SphK avoid apoptosis and thus continue to proliferate. By inhibiting SphK activity in the cancer cells, apoptosis is induced and the cancer

WO 01/60990

PCT/US01/04789

cells are destroyed. Essentially any cell proliferative disease or disorder that is dependent on SphK activity for cell survival or proliferation may be treated with an SphK activity inhibitor of the present invention. Examples include cell proliferation associated with vascular diseases, such as atherosclerosis or restenosis, with hyperplasia, or with cancers such as leukemia, lymphoma, ovarian cancer, breast cancer, lung cancer, colon cancer, testicular cancer, stomach cancer, or skin cancer. The cell proliferation may also be associated with angiogenesis. By preventing angiogenesis, the blood supply to a tumor may be diminished.

Therapeutics that antagonize activity may be administered in a therapeutic or prophylactic manner. Therapeutics that may be utilized include, but are not limited to: (i) an aforementioned peptide, or analogs, derivatives, fragments or homologs thereof; (ii) antibodies to an aforementioned peptide; (iii) nucleic acids encoding an aforementioned peptide; (iv) administration of antisense nucleic acid and nucleic acids that are "dysfunctional" (i.e., due to a heterologous insertion within the coding sequences of coding sequences to an aforementioned peptide) that are utilized to "knockout" endogenous function of an aforementioned peptide by homologous recombination (*see, e.g.*, Capecchi, 1989. *Science* 244: 1288-1292); or (v) modulators (i.e., inhibitors, agonists and antagonists, including additional peptide mimetic of the invention or antibodies specific to a peptide of the invention) that alter the interaction between an aforementioned peptide and its binding partner.

Diseases and disorders that are characterized by decreased (relative to a subject not suffering from the disease or disorder) levels or biological activity may be treated with therapeutics that increase (i.e., are agonists to) activity. This increase in activity may come about in a variety of ways, for example: (1) by increasing the copies of the gene in the cell (deamplifiers); (2) by increasing transcription of the SphK gene (transcription up-regulators); (3) by increasing the translation of SphK mRNA into protein (translation up-regulators); or (4) by increasing the activity of SphK itself (agonists).

In certain cells, decreased levels of SphK leads to apoptosis. In situations wherein cell survival or proliferation is desired, therapeutics that increase SphK activity are particularly useful. By increasing SphK activity in a cell, apoptosis is prevented. Cells that would otherwise be apoptotic are able to survive or proliferate. An example of a disease or disorder that may be treated with a therapeutic that increases SphK activity is ischemia, such as that associated with myocardial infarction. By promoting angiogenesis, the therapeutic's effect provides blood flow to the ischemic area. The therapeutics may also be used to increase peripheral blood flow to a specific area (e.g., a wound or a transplanted tissue, organ, limb).

Therapeutics that upregulate activity may be administered in a therapeutic or prophylactic manner. Therapeutics that may be utilized include, but are not limited to, an

WO 01/60990

PCT/US01/04789

aforementioned peptide, or analogs, derivatives, fragments or homologs thereof; or an agonist that increases bioavailability.

Increased or decreased levels can be readily detected by quantifying peptide and/or RNA, by obtaining a patient tissue sample (e.g., from biopsy tissue) and assaying it *in vitro* for RNA or peptide levels, structure and/or activity of the expressed peptides (or mRNAs of an
5 aforementioned peptide). Methods that are well-known within the art include, but are not limited to, immunoassays (e.g., by Western blot analysis, immunoprecipitation followed by sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis, immunocytochemistry, etc.) and/or hybridization assays to detect expression of mRNAs (e.g., Northern assays, dot blots, *in*
10 *situ* hybridization, and the like).

Prophylactic Methods

In one aspect, the invention provides a method for preventing, in a subject, a disease or condition associated with an aberrant SphK expression or activity, by administering to the subject an agent that modulates SphK expression or at least one SphK activity. Subjects at risk
15 for a disease that is caused or contributed to by aberrant SphK expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the SphK aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending upon the type of SphK aberrancy, for
20 example, an SphK agonist or SphK antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein. The prophylactic methods of the invention are further discussed in the following subsections.

Therapeutic Methods

Another aspect of the invention pertains to methods of modulating SphK expression or
25 activity for therapeutic purposes. The modulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of SphK protein activity associated with the cell. An agent that modulates SphK protein activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of an SphK protein, a peptide, an SphK peptidomimetic, or other small molecule. In one embodiment,
30 the agent stimulates one or more SphK protein activity. Examples of such stimulatory agents include active SphK protein and a nucleic acid molecule encoding SphK that has been introduced into the cell. In another embodiment, the agent inhibits one or more SphK protein activity. Examples of such inhibitory agents include antisense SphK nucleic acid molecules and anti-SphK antibodies. These modulatory methods can be performed *in vitro* (e.g., by culturing
35 the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject).

WO 01/60990

PCT/US01/04789

As such, the invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant expression or activity of an SphK protein or nucleic acid molecule. In one embodiment, the method involves administering an agent (*e.g.*, an agent identified by a screening assay described herein), or combination of agents that modulates (*e.g.*,
 5 up-regulates or down-regulates) SphK expression or activity. In another embodiment, the method involves administering an SphK protein or nucleic acid molecule as therapy to compensate for reduced or aberrant SphK expression or activity.

Inhibition of SphK activity is desirable in situations in which SphK is abnormally upregulated and/or in which decreased SphK activity is likely to have a beneficial effect (*e.g.*, to
 10 induce apoptosis). One example of such a situation is where a subject has a disorder characterized by aberrant cell proliferation and/or differentiation (*e.g.*, cancer or immune associated disorders). Another example of such a situation is where the subject has a gestational disease (*e.g.*, preclampsia).

Determination of the Biological Effect of the Therapeutic

15 In various embodiments of the invention, suitable *in vitro* or *in vivo* assays are performed to determine the effect of a specific therapeutic and whether its administration is indicated for treatment of the affected tissue.

In various specific embodiments, *in vitro* assays may be performed with representative cells of the type(s) involved in the patient's disorder, to determine if a given therapeutic exerts
 20 the desired effect upon the cell type(s). Compounds for use in therapy may be tested in suitable animal model systems including, but not limited to rats, mice, chicken, cows, monkeys, rabbits, and the like, prior to testing in human subjects. Similarly, for *in vivo* testing, any of the animal model system known in the art may be used prior to administration to human subjects.

Prophylactic and Therapeutic Uses of the Compositions of the Invention

25 The SphK nucleic acids and proteins of the invention are useful in potential prophylactic and therapeutic applications implicated in a variety of disorders including, but not limited to: proliferative disorders such as cancers, ischemia, or restenosis.

As an example, a cDNA encoding the SphK protein of the invention may be useful in gene therapy, and the protein may be useful when administered to a subject in need thereof. By
 30 way of non-limiting example, the compositions of the invention will have efficacy for treatment of patients suffering from cancers, ischemia, or restenosis.

Both the novel nucleic acid encoding the SphK protein, and the SphK protein of the invention, or fragments thereof, may also be useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed. A further use could be

WO 01/60990

PCT/US01/04789

as an anti-bacterial molecule (*i.e.*, some peptides have been found to possess anti-bacterial properties). These materials are further useful in the generation of antibodies that immunospecifically-bind to the novel substances of the invention for use in therapeutic or diagnostic methods.

5

EXAMPLES

The following examples illustrate by way of non-limiting example various aspects of the invention.

Example 1: Human Ortholog of Mouse AF068748

The inventor was able to piece together a homolog of the mouse SphK described in Kohama *et al.* (Kohama T, Olivera A, Edsall L, Nagiec MM, Dickson R, Spiegel S. "Molecular cloning and functional characterization of murine SphK". J Biol Chem. 1998 Sep 11;273(37):23722-8). Kohama *et al.* did not have conclusive data on where the mouse protein starts, even though they showed that the mouse cDNAs they obtained has SphKinase activity *in vivo*. However, no human sequence has been deposited for this enzyme in GenBank or Geneseq. In their paper, the authors have an alignment that shows part of a hypothetical human sequence. Here, the inventor obtained a virtual full-length human sequence with Kozak consensus.

Having the full-length human sequence permitted the inventor to establish where the start of the mouse protein occurs. The mouse homolog produced contains all the conserved regions the authors described in the paper.

Human SphK (ortholog of mouse AF068748)

This sequence was found while extending d010-107.9 (107.9) in job 6424 - collagen gel 24 hr versus 4, annotated as human similar to Norway rat gber_ai237625 EST234187. This led to Arabidopsis O65419 and C.Elegans T10B11.2, both proteins having some homology to mouse SphK AF068748. This sequence may be the human ortholog of AF068748 (because of the known involvement of this protein in protecting human umbilical cord vascular endothelial cells (HUVECs) from apoptosis (J Biol Chem, Vol. 274, Issue 48, 34499-34505, November 26, 1999). Similarities were sought using BlastN with this sequence against GenBankEST. Several human ESTs were identified that potentially code for the human ortholog: from 5' AI478197 (there is a repetitive sequence), D31133, AA026479, AA0232791, AA232646, AI769914, AI972156, and AA081152. These sequences were then capassembled in 2 groups, 5' (D31133, AA026479, AA0232791) and 3' (AI769914 and AA081152) groups and then the 2 groups were capassembled. In SEQCALLING™ 85667810 and 71311176 represent the 5' end. The capassembled sequence was fragmented. So the inventor seqextended using this sequence as

WO 01/60990

PCT/US01/04789

backbone and used the frameshift locations (that in general were weak assembly regions) to generate the right frame. The nucleic acid sequence of this human SphK cDNA is provided in SEQ ID NO:1. This sequence has a correct Kozak methionine. The amino acid sequence of the human SphK encoded by the cDNA is provided in SEQ ID NO:2.

5 Kohama *et al.* were concerned that the 2 ESTs they worked with had no Kozak methionine. All available mouse ESTs were seqextended and the consensus has a Kozak methionine (it starts at the same met and differs with SPHK1a by a couple of amino acids). The nucleic acid sequence of the consensus mouse SphK cDNA is provided by SEQ ID NO:3. The amino acid sequence of the mouse SphK encoded by the consensus cDNA is provided in SEQ
10 ID NO:4.

Example 2: Novel SphK

A novel gene fragment was differentially expressed using collagen gels at 24 hrs as compared to 4 hrs. AL096766 in HTG database Homo sapiens from clone 59H18 on chromosome 22 splicing 64154-64003,63017-62954,59856-59711,53578-53504,51333-
15 51181,47183-47078,45371-45295,43645-43440,42070-41860,39078-38779,37032-36151. In this clone there is a putative gene fragment annotated as dA59H18.2 (novel protein similar to human, mouse, yeast, worm and plant (predicted) proteins) created by splicing together 43440..43645,45295..45371, 47078..47183,51181..51333,53504..53578, 59711..59856, 62954..63017,64003..>64154). The isolation d010-107.9 (see Example 1) enabled the extension
20 of the gene in the 3' region, adding 93aa residues. In Seqcalling, the inventor assembled 85336019 (covers bases 915 to 1491 and 1800 to end), 84585492 (bases 360 to 960) and 80432911 that encodes for the 5' region and start codon. The cDNA was constructed by splicing 64154-64003,63017-62954,59856-59711,53578-53504,51333-51181,47183-47078,45371-45295,43645-43440,42070-41860,39078-38779,37032-36151, editing by
25 removing 37032 to 36151, and adding 80432911. The resulting nucleic acid sequence of this human SphK cDNA is provided in SEQ ID NO:5. The amino acid sequence of the human SphK encoded by this cDNA is provided in SEQ ID NO:6. This sequence extends the sequence deposited in GenBank as putative fragment peptide CAB62977 both at the NH and COOH termini for a total of 125aa (FIG. 1). The additions identify the actual start and add a further
30 conserved region (C5) to the deposited sequence.

A rat ortholog, designated AI237625, of this human SphK was determined. This ortholog is encoded in EST234187 from normalized rat placenta, Bento Soares Rattus sp. cDNA clone RPLDB60 3' end, mRNA sequence. A nucleic acid sequence comprising a portion of the

WO 01/60990

PCT/US01/04789

The choice of nucleic acid starting material, clone of interest, or library type is believed to be a matter of routine for a person of ordinary skill in the art with knowledge of the embodiments described herein. Other aspects, advantages, and modifications considered to be within the scope of the following claims.

WHAT IS CLAIMED IS:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:
 - (a) a mature form of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 6;
 - (b) a variant of a mature form of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 6, wherein one or more amino acid residues in said variant differs from the amino acid sequence of said mature form, provided that said variant differs in no more than 15% of the amino acid residues from the amino acid sequence of said mature form;
 - (c) an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 6; and
 - (d) a variant of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 6, wherein one or more amino acid residues in said variant differs from the amino acid sequence of said mature form, provided that said variant differs in no more than 15% of amino acid residues from said amino acid sequence.
2. The polypeptide of claim 1, wherein said polypeptide comprises the amino acid sequence of a naturally-occurring allelic variant of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 6.
3. The polypeptide of claim 2, wherein said allelic variant comprises an amino acid sequence that is the translation of a nucleic acid sequence differing by a single nucleotide from a nucleic acid sequence selected from the group consisting of SEQ ID NOS:1 and 5.
4. The polypeptide of claim 1, wherein the amino acid sequence of said variant comprises a conservative amino acid substitution.
5. An isolated nucleic acid molecule comprising a nucleic acid sequence encoding a polypeptide comprising an amino acid sequence selected from the group consisting of:
 - (a) a mature form of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 6;
 - (b) a variant of a mature form of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 6, wherein one or more amino acid residues in

WO 01/60990

PCT/US01/04789

- said variant differs from the amino acid sequence of said mature form, provided that said variant differs in no more than 15% of the amino acid residues from the amino acid sequence of said mature form;
- (c) an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 6;
 - (d) a variant of an amino acid sequence selected from the group consisting of SEQ ID NOS:2 and 6, wherein one or more amino acid residues in said variant differs from the amino acid sequence of said mature form, provided that said variant differs in no more than 15% of amino acid residues from said amino acid sequence;
 - (e) a nucleic acid fragment encoding at least a portion of a polypeptide comprising an amino acid sequence chosen from the group consisting of SEQ ID NOS:2 and 6, or a variant of said polypeptide, wherein one or more amino acid residues in said variant differs from the amino acid sequence of said mature form, provided that said variant differs in no more than 15% of amino acid residues from said amino acid sequence; and
 - (f) a nucleic acid molecule comprising the complement of (a), (b), (c), (d) or (e).
6. The nucleic acid molecule of claim 5, wherein the nucleic acid molecule comprises the nucleotide sequence of a naturally-occurring allelic nucleic acid variant.
 7. The nucleic acid molecule of claim 5, wherein the nucleic acid molecule encodes a polypeptide comprising the amino acid sequence of a naturally-occurring polypeptide variant.
 8. The nucleic acid molecule of claim 5, wherein the nucleic acid molecule differs by a single nucleotide from a nucleic acid sequence selected from the group consisting of SEQ ID NOS:1 and 5.
 9. The nucleic acid molecule of claim 5, wherein said nucleic acid molecule comprises a nucleotide sequence selected from the group consisting of:
 - (a) a nucleotide sequence selected from the group consisting of SEQ ID NOS:1 and 5;
 - (b) a nucleotide sequence differing by one or more nucleotides from a nucleotide sequence selected from the group consisting of SEQ ID NOS:1 and 5, provided that no more than 20% of the nucleotides differ from said nucleotide sequence;
 - (c) a nucleic acid fragment of (a); and

WO 01/60990

PCT/US01/04789

- (d) a nucleic acid fragment of (b).
- 10. The nucleic acid molecule of claim 5, wherein said nucleic acid molecule hybridizes under stringent conditions to a nucleotide sequence chosen from the group consisting of SEQ ID NOS:1 and 5, or a complement of said nucleotide sequence.
- 11. The nucleic acid molecule of claim 5, wherein the nucleic acid molecule comprises a nucleotide sequence selected from the group consisting of:
 - (a) a first nucleotide sequence comprising a coding sequence differing by one or more nucleotide sequences from a coding sequence encoding said amino acid sequence, provided that no more than 20% of the nucleotides in the coding sequence in said first nucleotide sequence differ from said coding sequence;
 - (b) an isolated second polynucleotide that is a complement of the first polynucleotide; and
 - (c) a nucleic acid fragment of (a) or (b).
- 12. A vector comprising the nucleic acid molecule of claim 11.
- 13. The vector of claim 12, further comprising a promoter operably-linked to said nucleic acid molecule.
- 14. A cell comprising the vector of claim 12.
- 15. An antibody that binds immunospecifically to the polypeptide of claim 1.
- 16. The antibody of claim 15, wherein said antibody is a monoclonal antibody.
- 17. The antibody of claim 15, wherein the antibody is a humanized antibody.
- 18. A method for determining the presence or amount of the polypeptide of claim 1 in a sample, the method comprising:
 - (a) providing the sample;
 - (b) contacting the sample with an antibody that binds immunospecifically to the polypeptide; and
 - (c) determining the presence or amount of antibody bound to said polypeptide, thereby determining the presence or amount of polypeptide in said sample.
- 19. A method for determining the presence or amount of the nucleic acid molecule of claim 5 in a sample, the method comprising:

WO 01/60990

PCT/US01/04789

- (a) providing the sample;
 - (b) contacting the sample with a probe that binds to said nucleic acid molecule; and
 - (c) determining the presence or amount of the probe bound to said nucleic acid molecule,
- thereby determining the presence or amount of the nucleic acid molecule in said sample.
20. The method of claim 19 wherein presence or amount of the nucleic acid molecule is used as a marker for cell or tissue type.
21. The method of claim 20 wherein the cell or tissue type is cancerous.
22. A method of identifying an agent that binds to a polypeptide of claim 1, the method comprising:
- (a) contacting said polypeptide with said agent; and
 - (b) determining whether said agent binds to said polypeptide.
23. The method of claim 22 wherein the agent is a cellular receptor or a downstream effector.
24. A method for identifying an agent that modulates the expression or activity of the polypeptide of claim 1, the method comprising:
- (a) providing a cell expressing said polypeptide;
 - (b) contacting the cell with said agent, and
 - (c) determining whether the agent modulates expression or activity of said polypeptide,
- whereby an alteration in expression or activity of said peptide indicates said agent modulates expression or activity of said polypeptide.
25. A method for modulating the activity of the polypeptide of claim 1, the method comprising contacting a cell sample expressing the polypeptide of said claim with a compound that binds to said polypeptide in an amount sufficient to modulate the activity of the polypeptide.
26. A method of treating or preventing a SphK-associated disorder, said method comprising administering to a subject in which such treatment or prevention is desired the polypeptide of claim 1 in an amount sufficient to treat or prevent said SphK-associated disorder in said subject.

- 80

WO 01/60990

PCT/US01/04789

42. A kit comprising in one or more containers, the pharmaceutical composition of claim 39.
43. A kit comprising in one or more containers, the pharmaceutical composition of claim 40.
44. A method for determining the presence of or predisposition to a disease associated with altered levels of the polypeptide of claim 1 in a first mammalian subject, the method comprising:
 - (a) measuring the level of expression of the polypeptide in a sample from the first mammalian subject; and
 - (b) comparing the amount of said polypeptide in the sample of step (a) to the amount of the polypeptide present in a control sample from a second mammalian subject known not to have, or not to be predisposed to, said disease;
wherein an alteration in the expression level of the polypeptide in the first subject as compared to the control sample indicates the presence of or predisposition to said disease.
45. The method of claim 44 wherein the predisposition is to cancers.
46. A method for determining the presence of or predisposition to a disease associated with altered levels of the nucleic acid molecule of claim 5 in a first mammalian subject, the method comprising:
 - (a) measuring the amount of the nucleic acid in a sample from the first mammalian subject; and
 - (b) comparing the amount of said nucleic acid in the sample of step (a) to the amount of the nucleic acid present in a control sample from a second mammalian subject known not to have or not be predisposed to, the disease;
wherein an alteration in the level of the nucleic acid in the first subject as compared to the control sample indicates the presence of or predisposition to the disease.
47. The method of claim 46 wherein the predisposition is to cancers.
48. A method of treating a pathological state in a mammal, the method comprising administering to the mammal a polypeptide in an amount that is sufficient to alleviate the pathological state, wherein the polypeptide is a polypeptide having an amino acid sequence at least 95% identical to a polypeptide comprising an amino acid sequence of at least one of SEQ ID NOS:2 and 6, or a biologically active fragment thereof.

WO 01/60990

PCT/US01/04789

49. A method of treating a pathological state in a mammal, the method comprising administering to the mammal an isolated nucleic acid in an amount that is sufficient to alleviate the pathological state, wherein the isolated nucleic acid is an isolated nucleic acid having an nculeic acid sequence at least 95% identical to a nucleic acid comprising a nucleic acid sequence of at least one of SEQ ID NOS:2 and 6, or a compliment thereof.
50. A method of treating a pathological state in a mammal, the method comprising administering to the mammal the antibody of claim 15 in an amount sufficient to alleviate the pathological state.
51. A non-human transgenic animal comprising a transgene comprising a nucleic acid of claim 5.
52. A method for identifying an agent that modulates the expression or activity of the polypeptide of claim 1, the method comprising:
 - (a) contacting a non-human transgenic animal of claim 51 with said agent; and
 - (b) determining whether the agent modulates expression or activity of said polypeptide,whereby an alteration in expression or activity of said peptide indicates said agent modulates expression or activity of said polypeptide.
53. A non-human transgenic animal comprising a transgene disrupting an endogenous sphingosine kinase gene.

FIG. 1

Multiple Alignment:	
Multiple Alignment:	
SP1	MEIPYAFTVHCVKRAERRHWKWAQWTFWCPEEQCLHLWLQTLREMLEKLTSEPKHLLVFI
CA	SPkinaseinHSDA59H18
SP1	PKHLLVFI
CA	CAB62977
SP1	NPFGGKGQGGKEIYERKVAPLFTLASITTDIIGNKEFYVN YVEVITEHANQAKETLYEINID
CA	SPkinaseinHSDA59H18
SP1	NPFGGKGQGGKEIYERKVAPLFTLASITTDIIGNKEFYVN YVEVITEHANQAKETLYEINID
CA	CAB62977
SP1	KYDGINVCVGGDGMFSEVLHGLIGRTQSSAGVDQNHPRALVLPSSLRIGIIPAGSTDCVCY
CA	SPkinaseinHSDA59H18
SP1	KYDGINVCVGGDGMFSEVLHGLIGRTQSSAGVDQNHPRALVLPSSLRIGIIPAGSTDCVCY
CA	CAB62977
SP1	STVGTSDAETSALHI VVGDSLAMDVSS VHHNSTLLREYS VSLGCGFYGDIINDSEKKRWL
CA	SPkinaseinHSDA59H18
SP1	STVGTSDAETSALHI VVGDSLAMDVSS VHHNSTLLREYS VSLGCGFYGDIINDSEKKRWL
CA	CAB62977
SP1	GLAEYDFSGLIKTF LSHHCYEGT VSF LPAQHTVGS PEDRKPCRAGCFVCRQSKQQLEEEQK
CA	SPkinaseinHSDA59H18
SP1	GLAEYDFSGLIKTF LSHHCYEGT VSF LPAQHTVGS PEDRKPCRAGCFVCRQSKQQLEEEQK
CA	CAB62977
SP1	KALYGLEAAEDVEEWQVVGKFLAINATHMSCACRRSP RGLSPAHLGDGSSDLILIRKC
CA	SPkinaseinHSDA59H18
SP1	KALYGLEAAEDVEEWQVVGKFLAINATHMSCACRRSP RGLSPAHLGDGSSDLILIRKC
CA	CAB62977
SP1	SEFNFLRFLIRHTNQDQDFTFVEVYRVKKEFQFTSKHMEDESDLEKGGKRFQGHICSS
CA	SPkinaseinHSDA59H18
SP1	SEFNFLRFLIRHTNQDQD
CA	CAB62977
SP1	HPSCCCTVSNSSWNCDGEVLHSPAIEVEVHCQLVELFA RGLIENPIEPDSHS
CA	SPkinaseinHSDA59H18
SP1	
CA	CAB62977

WO 01/60990

PCT/US01/04789

FIG. 2

Multiple Alignment:

```

52180312 -----
A1237625 -----
SPkinaseinHSDA59H18 PKHLLVF INPFGGKGQKRIYERKVAPLFTLASITTDIGNKFVNVYVEVITEHANQAKE
-----
52180312 -----
A1237625 -----
SPkinaseinHSDA59H18 TLYEINI DKYDGI VCVGGDMFSEVLHGLIGRTQRSAGVDQNHPRAVLPSSLRIGIIPA
-----
52180312 -----
A1237625 -----
SPkinaseinHSDA59H18 GSTDCVCYSTVGTSDAETSA LHI VVGDSLAMDVSSVHHNSTLLRYSVSLG YGFYGDIIK
-----
52180312 -----
A1237625 -----
SPkinaseinHSDA59H18 DSEKKRWLGLARYDFSGLK TFLSHHCYEGTVSFLPAQHTVGS PRDRKPCRAGCFVCRQSK
-----
52180312 -----
A1237625 -----
SPkinaseinHSDA59H18 QQL EEEQKKALYGLEAAEDVEEWQVVC GKFLAINATNMSCACRSPRFLSPAHLGDGSS
-----
52180312 -----
A1237625 -----
SPkinaseinHSDA59H18 LLLILIEKCSRNFNLFRLIEHTNQEDQFGFTFVEVYRWKRFQFTSEHVELDNDLKELEK
-----
52180312 -----
A1237625 -----
SPkinaseinHSDA59H18 LLLILIEKCSRNFNLFRLIEHTNQEDQFDFTFVEVYRWKRFQFTSEHVELDNDLKELEK
-----
52180312 -----
A1237625 -----
SPkinaseinHSDA59H18 KFGKICIEDRPSCTPSASRSSWNC DGEVWHSP AIEVEVWHCQLVELFARGIEEES
-----
52180312 -----
A1237625 -----
SPkinaseinHSDA59H18 KFGQICIEDNPPPAAPTSSSSWNC DGEVLHSP AIEVEVWHCQLVELFARGIEEES
-----
52180312 -----
A1237625 -----
SPkinaseinHSDA59H18 KFGHICSSHPSCCEIIVENSSWNC DGEVLHSP AIEVEVWHCQLVELFARGIEENPKPDSHS

```


WO 01/60990

PCT/US01/04789

FIG. 4

PHYLIP - Protein Distance Analysis

Sequences analyzed:


1. SPkinaseinHSDA59H18
2. Q18425
3. O14159
4. Q06147
5. Q12246
6. coorrected_human_sphingosine
7. AF068748_EXT-2

7 Populations

Neighbor-Joining/UPGMA method version 3.572c

Neighbor-joining method

Negative branch lengths allowed



remember: this is an unrooted tree!

Between	And	Length
5	3	1.09970
3	1	1.59865
1	Q06147	0.79847
1	Q12246	1.02742
3	O14159	1.74337
5	Q18425	4.00993
5	4	0.43259
4	2	1.44382
2	coorrected_	0.24774
2	AF068748_EXT	0.37392
4	SPkinasein	3.26442

WO 01/60990

PCT/US01/04789

FIG. 5

Multiple Alignment:

80432911 SPKinsaiHSDA59H18	AGAPCALACGPTVSEIIAVEETVHGKHGGSGKHWQIMETVYFTVHGVKFAREHFWKWAC -----MEIPVAFTVHGVKFAREHFWKWAC
80432911 SPKinsaiHSDA59H18	VTFWCPPEEQCLHLWLQTLREMLEILLTSEPHELLVFINPFGSGGGGIEIYERKVAFLFTLA VTFWCPPEEQCLHLWLQTLREMLEILLTSEPHELLVFINPFGSGGGGIEIYERKVAFLFTLA
80432911 SPKinsaiHSDA59H18	SITTDII-----TEHANQAKETLVEINIDKYVSGIWCWGGSDGMPFSEVLHGLIGR SITTDIIQHNFYVNVYEWITEHANQAKETLVEINIDKYVSGIWCWGGSDGMPFSEVLHGLIGR
80432911 SPKinsaiHSDA59H18	TQESAGVDQNHPR-----TQESAGVLCQNHPEA VLWPEELRIGIIPAGSTGVCYSTGTSDAETSAHLIWWGDSLAMC TQESAGVLCQNHPEA VLWPEELRIGIIPAGSTGVCYSTGTSDAETSAHLIWWGDSLAMC
80432911 SPKinsaiHSDA59H18	WSSWHHNSLTLETSVLLGGVGTGGIHDSEHNRWLGLARTDFGRLETPLCHHCYEYGVG -----
80432911 SPKinsaiHSDA59H18	FLPAQHTWGGPRDREHFRAGCPWRQGGTQQLLEEQKIALVGLAAEDVEEWQWNGGIFLA -----
80432911 SPKinsaiHSDA59H18	IIAATHMSCACERSPEGLSPAHLGGSSDLIIKESSEFNFLERLTHHQDQDQDFTFF -----
80432911 SPKinsaiHSDA59H18	EYVRVLIIFQTSIEHMEDEGLDIEGGHFRGHICGCHPSGGCTVSGSWIWDGCVLHCPA -----
80432911 SPKinsaiHSDA59H18	IEVRVHCQLVELFARCTIENPIEDSHC -----

WO 01/60990

PCT/US01/04789

```

gcccggtggtc gccttccgct tggagcccaa ggatgggaaa ggtgtgtttg cagtggatgg 1200
ggaattgatg gtttagcgagg ccgtgcaggg ccaggtgcac ccaaactact tctggatggt 1260
cagcgggttg gtggagcccc cgcccagctg gaagcccag cagatgccac cgccagaaga 1320
gcccttatga cccctgggccc gcgtgtgtcc ttagtgtcta cttgcaggac ccttcctcct 1380
tccctagggc tgcagggcct gtccacagct cctgtggggg tggaggagac tcctctggag 1440
aaggggtgaga aggtggaggc tatgctttgg ggggacaggc cagaatgaag tcctgggtca 1500
ggagcccagc tggtggggc cagctgccta tgtaaggcct tctagtttgt tctgagacct 1560
ccacccacag aaccaaattc aaataaagtg acattcccaa 1600

```

<210> 2

<211> 384

<212> PRT

<213> Homo sapiens

<400> 2

```

Met Asp Pro Ala Gly Gly Pro Arg Gly Val Leu Pro Arg Pro Cys Arg
 1           5           10           15

```

```

Val Leu Val Leu Leu Asn Pro Arg Gly Gly Lys Gly Lys Ala Leu Gln
          20           25           30

```

```

Leu Phe Arg Ser His Val Gln Pro Leu Leu Ala Glu Ala Glu Ile Ser
          35           40           45

```

```

Phe Thr Leu Met Leu Thr Glu Arg Arg Asn His Ala Arg Glu Leu Val
          50           55           60

```

```

Arg Ser Glu Glu Leu Gly Arg Trp Asp Ala Leu Val Val Met Ser Gly
          65           70           75           80

```

```

Asp Gly Leu Met His Glu Val Val Asn Gly Leu Met Glu Arg Pro Asp
          85           90           95

```

```

Trp Glu Thr Ala Ile Gln Lys Pro Leu Cys Ser Leu Pro Ala Gly Ser
          100          105          110

```

```

Gly Asn Ala Leu Ala Ala Ser Leu Asn His Tyr Ala Gly Tyr Glu Gln
          115          120          125

```

```

Val Thr Asn Glu Asp Leu Leu Thr Asn Cys Thr Leu Leu Leu Cys Arg
          130          135          140

```

```

Pro Val Leu Ser Pro Met Asn Leu Leu Ser Leu His Thr Ala Ser Gly
          145          150          155          160

```

```

Leu Arg Ser Phe Ser Val Leu Ser Leu Ala Trp Gly Phe Ile Ala Asp
          165          170          175

```


WO-01/60990

PCT/US01/04789

Val Asp Leu Glu Ser Asp Lys Tyr Arg Arg Leu Gly Glu Met Arg Phe
180 185 190

Thr Leu Gly Thr Phe Leu Arg Leu Ala Ala Leu Arg Thr Tyr Arg Gly
195 200 205

Arg Leu Ala Thr Leu Pro Val Gly Arg Val Gly Phe Lys Thr Pro Ala
210 215 220

Ser Pro Val Val Val Gln Gln Gly Pro Val Asp Ala His Leu Val Pro
225 230 235 240

Leu Glu Glu Gln Val Pro Ser His Trp Gln Val Val Pro Asp Glu Asp
245 250 255

Phe Val Leu Val Leu Ala Leu Leu His Ser His Leu Ala Ser Glu Met
260 265 270

Phe Ala Ala Pro Met Gly Arg Cys Ala Ala Gly Val Met His Leu Phe
275 280 285

Tyr Val Arg Ala Gly Val Ser Arg Ala Met Leu Leu Arg Leu Phe Leu
290 295 300

Ala Met Glu Lys Gly Arg His Met Glu Tyr Glu Cys Pro Tyr Leu Val
305 310 315 320

Tyr Val Pro Val Val Ala Phe Arg Leu Glu Pro Lys Asp Gly Lys Gly
325 330 335

Val Phe Ala Val Asp Gly Glu Leu Met Val Ser Glu Ala Val Gln Gly
340 345 350

Gln Val His Pro Asn Tyr Phe Trp Met Val Ser Gly Cys Val Glu Pro
355 360 365

Pro Pro Ser Trp Lys Pro Gln Gln Met Pro Pro Pro Glu Glu Pro Leu
370 375 380

<210> 3

<211> 1759

<212> DNA

<213> Mus musculus

WO 01/60990

PCT/US01/04789

<400> 3

```

gaactccagg gtcctctggg agagaagcca cttcaaggc gtgacctagt tcaactgcaat 60
cctttcttat ctgggttcgt ttctctcttg gactcgctc ttctggactt taagaagcga 120
tgcgaagata gagatctggc cgtcccgagg aatgacgagg gcgtcacac agcccaggga 180
ggtggagagg gcgagcccac ggccagtcgc cagacaccct cctgggcaac accgataaga 240
agctgaacgc aggagccgcc gttacctcta gcagcgccgg ggagcaccg gtggccctt 300
gtcagcgga gcccgggac ctggtatgg aaccagtaga atgccctcga ggactgctcc 360
cacggccatg cagagtgtg gtgtgtgta accccaggg tggcaagggc aaggctctgc 420
agctcttcca gagccgtgtg cagcccttcc tggaggagg agagataacc tttaaactga 480
tactcacga acggaagaac catgccagg agctggtgtg tgcagaggag ttgggtcact 540
gggagccct ggagtcctg tccggtgat gtctgatgca tgaagtgtg aatgggcta 600
tggaacggc agactgggag actgccatcc agaaaccct gtgtagctc cctggaggct 660
ccggcaatgc gctggcagct tctgtgaacc actatgctg gtacgagcag gtgactaatg 720
aagacctgct catcaactgc aactgctgt tgtgcgcgc gcgcctgtca cccatgaacc 780
tgctgtccct gcacactgct tctgggctgc ggtctctattc tgtgtcagt ctgtcctggg 840
gctttgttgc tgacgtggac ctgagagtg agaagtacag gcgttgggg gagattcgtt 900
tcacagtggg cacttcttt cgcctagcaa gcctgcgcct ctaccaaggc caactggcct 960
accttctgt aggaactgtg gcctctaaga gaccgcctc tacactggtg cagaagggcc 1020
ccgtcgacac acacttgtt cctctggagg agccagtgc ttctcattgg actgtgttac 1080
cagaacagga cttcgtctc gtgtgtgtg tgctacacac ccacctgagc tccgagctgt 1140
ttgcagcacc catgggcgc tgtgaggct gtgttatgca tctgttctac gtacgtgcg 1200
gggtgtcaag ggctgcgct ctgcgcctc tcctggccat gcagaagggc aagcatatg 1260
aacttgactg tccatacctg gttcatgtg ccgtggttg tttccgcctg gagcccagga 1320
gccagagggg cgtgttttct gtggatggg agctgatgt atgtgaagct gtgcagggcc 1380
aagtgcaccc aaactacct tggatggtc gtggcagcag agatgcccc tccggccggg 1440
actcccgccg ggggccacct ccagaagaac cataactctg tgcctttgt tactctgtc 1500
aggctgagat gggaccctc cccgcacca cctcctggt tgggagggt tttctaaagt 1560
tcctatggaa gtgtgggga cccctgcga gaaagctaga aggtggggt atgacttgga 1620
aagaaaggct ttaccttcca gttagagtaa catccccagt agagccctgc tggctggacc 1680
agttgcatat agaagacatt cccattgct tttagggacc ttccctggga accaaattca 1740
aataaagaga cttttccaa 1759

```

<210> 4

<211> 382

<212> PRT

<213> Mus musculus

<400> 4

```

Met Glu Pro Val Glu Cys Pro Arg Gly Leu Leu Pro Arg Pro Cys Arg
 1             5             10             15

Val Leu Val Leu Leu Asn Pro Gln Gly Gly Lys Gly Lys Ala Leu Gln
      20             25             30

Leu Phe Gln Ser Arg Val Gln Pro Phe Leu Glu Glu Ala Glu Ile Thr
 35             40             45

```

WO 01/60990

PCT/US01/04789

Phe Lys Leu Ile Leu Thr Glu Arg Lys Asn His Ala Arg Glu Leu Val
50 55 60

Cys Ala Glu Glu Leu Gly His Trp Asp Ala Leu Ala Val Met Ser Gly
65 70 75 80

Asp Gly Leu Met His Glu Val Val Asn Gly Leu Met Glu Arg Pro Asp
85 90 95

Trp Glu Thr Ala Ile Gln Lys Pro Leu Cys Ser Leu Pro Gly Gly Ser
100 105 110

Gly Asn Ala Leu Ala Ala Ser Val Asn His Tyr Ala Gly Tyr Glu Gln
115 120 125

Val Thr Asn Glu Asp Leu Leu Ile Asn Cys Thr Leu Leu Leu Cys Arg
130 135 140

Arg Arg Leu Ser Pro Met Asn Leu Leu Ser Leu His Thr Ala Ser Gly
145 150 155 160

Leu Arg Leu Tyr Ser Val Leu Ser Leu Ser Trp Gly Phe Val Ala Asp
165 170 175

Val Asp Leu Glu Ser Glu Lys Tyr Arg Arg Leu Gly Glu Ile Arg Phe
180 185 190

Thr Val Gly Thr Phe Phe Arg Leu Ala Ser Leu Arg Ile Tyr Gln Gly
195 200 205

Gln Leu Ala Tyr Leu Pro Val Gly Thr Val Ala Ser Lys Arg Pro Ala
210 215 220

Ser Thr Leu Val Gln Lys Gly Pro Val Asp Thr His Leu Val Pro Leu
225 230 235 240

Glu Glu Pro Val Pro Ser His Trp Thr Val Val Pro Glu Gln Asp Phe
245 250 255

Val Leu Val Leu Val Leu Leu His Thr His Leu Ser Ser Glu Leu Phe
260 265 270

Ala Ala Pro Met Gly Arg Cys Glu Ala Gly Val Met His Leu Phe Tyr
275 280 285

Val Arg Ala Gly Val Ser Arg Ala Ala Leu Leu Arg Leu Phe Leu Ala
290 295 300

WO 01/60990

PCT/US01/04789

Met Gln Lys Gly Lys His Met Glu Leu Asp Cys Pro Tyr Leu Val His
305 310 315 320

Val Pro Val Val Ala Phe Arg Leu Glu Pro Arg Ser Gln Arg Gly Val
325 330 335

Phe Ser Val Asp Gly Glu Leu Met Val Cys Glu Ala Val Gln Gly Gln
340 345 350

Val His Pro Asn Tyr Leu Trp Met Val Cys Gly Ser Arg Asp Ala Pro
355 360 365

Ser Gly Arg Asp Ser Arg Arg Gly Pro Pro Pro Glu Glu Pro
370 375 380

<210> 5

<211> 1840

<212> DNA

<213> Homo sapiens

<400> 5

gccggcgccc ccggcgcgga tgctgtctct gtgcctgtat ctgagatcat cgccgttgag 60
gaaacagacg ttcacgggaa acatcaaggc agtggaaaat ggcagaaaat ggaaaagcct 120
tacgctttta cagttcactg tgtaaagaga gcacgacggc accgctggaa gtgggcgcag 180
gtgactttct ggtgtccaga ggagcagctg tgtcacttgt ggctgcagac cctgcgggag 240
atgctggaga agctgacgtc cagaccaaag catttacttg tatttatcaa cccgtttgga 300
ggaaaaggac aaggcaagcg gatatatgaa agaaaagtgg caccactgtt caccttagcc 360
tccatcacca ctgacatcat cggtaacaaa ttctatgtta actatgtaga agtaattact 420
gaacatgcta atcaggccaa ggagactctg tatgagatta acatagacaa atacgacggc 480
atcgtctgtg tcggcgagga tggatgttc agcgaggtgc tgcacgttct gattgggagg 540
acgcagagga gcgcgggggt cgaccagaac ccccccggt ctgtgtctgt cccagtagc 600
ctccggattg gaatcattcc cgcagggtca acggactgcg tgtgttactc caccgtgggc 660
accagcgacg cagaaacctc ggcgctgcat atcgttgttg gggactcgtt ggccatggat 720
gtgtctctag tccaccacaa cagcacactc ctctgtactt cctgttccct gctgggctac 780
ggcttctacg gggacatcat caaggacagt gagaagaaac ggtggttggg tcttgccaga 840
tacgactttt caggtttaaa gaccttctc tcccaccact gctatgaagg gacagtgtcc 900
ttctctcctg cacaacacac ggtgggatct ccaagggata ggaagccctg ccgggcagga 960
tgctttgttt gcaggcaaag caagcagcag ctggaggagg agcagaagaa agcactgtat 1020
ggtttggaag ctgcggagga cgtggaggag tggcaagtgc tctgtgggaa gtttctggcc 1080
atcaatgcca caaacatgtc ctgtgcttgt cgccggagcc ccaggggcct ctccccggt 1140
gcccacttgg gagacgggtc ttctgacctc atctcatcc ggaaatgctc caggttcaat 1200
tttctgagat ttctcatcag gcacaccaac cagcaggacc agtttgactt cacttttgtt 1260
gaagtttatc gcgtcaagaa attccagttt acgtcgaagc acatggagga tgaggacagc 1320
gacctcaagg agggggggaa gaagcgcttt gggcacattt gcagcagcca cccctcctgc 1380
tgctgcaccg tctccaacag ctcttggaac tgcgacgggg aggtcctgca cagccctgcc 1440
atcgagggtca ggttccactg ccagctggtt cgactcttgc cagaggaat tgaagagaat 1500

WO 01/60990

PCT/US01/04789

ccgaagccag actcacacag ctgagaagcc ggcgtcctgc tcacaaactg ggaaagtgtg 1560
 aaaactatatt aagataatta ttacagacca attatgttga tatatacatt taaatgtaga 1620
 aatttatattt tgatagttaa atcttgattt tagaagaaaa cccttttgtc aacaattttg 1680
 tgtacatatt tggcattttc agttctgtac gcatctgcgg gttgcagccc acgccgctta 1740
 ctctcagcaa gaaagaggtg gaggttgagg tgagccaaga ttgcgccact gcaactccagc 1800
 ctggggcaaac agagggagac tccatcgccc cccccaacaa 1840

<210> 6

<211> 471

<212> PRT

<213> Homo sapiens

<400> 6

Met Glu Lys Pro Tyr Ala Phe Thr Val His Cys Val Lys Arg Ala Arg
 1 5 10 15

Arg His Arg Trp Lys Trp Ala Gln Val Thr Phe Trp Cys Pro Glu Glu
 20 25 30

Gln Leu Cys His Leu Trp Leu Gln Thr Leu Arg Glu Met Leu Glu Lys
 35 40 45

Leu Thr Ser Arg Pro Lys His Leu Leu Val Phe Ile Asn Pro Phe Gly
 50 55 60

Gly Lys Gly Gln Gly Lys Arg Ile Tyr Glu Arg Lys Val Ala Pro Leu
 65 70 75 80

Phe Thr Leu Ala Ser Ile Thr Thr Asp Ile Ile Gly Asn Lys Phe Tyr
 85 90 95

Val Asn Tyr Val Glu Val Ile Thr Glu His Ala Asn Gln Ala Lys Glu
 100 105 110

Thr Leu Tyr Glu Ile Asn Ile Asp Lys Tyr Asp Gly Ile Val Cys Val
 115 120 125

Gly Gly Asp Gly Met Phe Ser Glu Val Leu His Gly Leu Ile Gly Arg
 130 135 140

Thr Gln Arg Ser Ala Gly Val Asp Gln Asn His Pro Arg Ala Val Leu
 145 150 155 160

Val Pro Ser Ser Leu Arg Ile Gly Ile Ile Pro Ala Gly Ser Thr Asp
 165 170 175

Cys Val Cys Tyr Ser Thr Val Gly Thr Ser Asp Ala Glu Thr Ser Ala

WO 01/60990

PCT/US01/04789

180	185	190
Leu His Ile Val Val Gly Asp Ser Leu Ala Met Asp Val Ser Ser Val		
195	200	205
His His Asn Ser Thr Leu Leu Arg Tyr Ser Val Ser Leu Leu Gly Tyr		
210	215	220
Gly Phe Tyr Gly Asp Ile Ile Lys Asp Ser Glu Lys Lys Arg Trp Leu		
225	230	235
Gly Leu Ala Arg Tyr Asp Phe Ser Gly Leu Lys Thr Phe Leu Ser His		
245	250	255
His Cys Tyr Glu Gly Thr Val Ser Phe Leu Pro Ala Gln His Thr Val		
260	265	270
Gly Ser Pro Arg Asp Arg Lys Pro Cys Arg Ala Gly Cys Phe Val Cys		
275	280	285
Arg Gln Ser Lys Gln Gln Leu Glu Glu Glu Gln Lys Lys Ala Leu Tyr		
290	295	300
Gly Leu Glu Ala Ala Glu Asp Val Glu Glu Trp Gln Val Val Cys Gly		
305	310	315
Lys Phe Leu Ala Ile Asn Ala Thr Asn Met Ser Cys Ala Cys Arg Arg		
325	330	335
Ser Pro Arg Gly Leu Ser Pro Ala Ala His Leu Gly Asp Gly Ser Ser		
340	345	350
Asp Leu Ile Leu Ile Arg Lys Cys Ser Arg Phe Asn Phe Leu Arg Phe		
355	360	365
Leu Ile Arg His Thr Asn Gln Gln Asp Gln Phe Asp Phe Thr Phe Val		
370	375	380
Glu Val Tyr Arg Val Lys Lys Phe Gln Phe Thr Ser Lys His Met Glu		
385	390	395
Asp Glu Asp Ser Asp Leu Lys Glu Gly Gly Lys Lys Arg Phe Gly His		
405	410	415
Ile Cys Ser Ser His Pro Ser Cys Cys Cys Thr Val Ser Asn Ser Ser		
420	425	430
Trp Asn Cys Asp Gly Glu Val Leu His Ser Pro Ala Ile Glu Val Arg		

WO 01/60990

PCT/US01/04789

435

440

445

Val His Cys Gln Leu Val Arg Leu Phe Ala Arg Gly Ile Glu Glu Asn
450 455 460

Pro Lys Pro Asp Ser His Ser
465 470

<210> 7

<211> 522

<212> DNA

<213> Rattus sp.

<400> 7

gtggatatgt caacataatt ggtcagtcga aattttctta agtaatatcc acacttccca 60
aagctcctgg gctttgggat tcttccttag gactcttctt cgatcccccg agcaaagagg 120
cgcaccagct ggtagtgac cctgacctca atggcagggc tgtgcaggac ttcccatca 180
cagttccagg agcttctaga ggttgggcag gcgcaagggg ggttgtcctt gcagatctgc 240
ccaaacttct gcttctctag ttccttcaag tcattgtcat catcttccac gtgctttgat 300
gtgaactgga atttcttgac tcgataaact tcaacgaaag tgaagccaaa ctgatcctcc 360
tggttggtgt gccggatgag gaatctcagg aagttgaacc tggagcattt ccggataagg 420
atgaggtcag aagacccatc tcccagatgg gcaaattggg acaggccccc agggctccgg 480
ggacaagcac aggacatgtt ggtggagttg atagccagga ac 522

<210> 8

<211> 144

<212> PRT

<213> Rattus sp.

<400> 8

Phe Leu Ala Ile Asn Ser Thr Asn Met Ser Cys Ala Cys Pro Arg Ser
1 5 10 15

Pro Gly Gly Leu Ser Pro Phe Ala His Leu Gly Asp Gly Ser Ser Asp
20 25 30

Leu Ile Leu Ile Arg Lys Cys Ser Arg Phe Asn Phe Leu Arg Phe Leu
35 40 45

Ile Arg His Thr Asn Gln Glu Asp Gln Phe Gly Phe Thr Phe Val Glu
50 55 60

Val Tyr Arg Val Lys Lys Phe Gln Phe Thr Ser Lys His Val Glu Asp
65 70 75 80

Asp Asp Asn Asp Leu Lys Glu Leu Glu Lys Gln Lys Phe Gly Gln Ile

WO 01/60990

PCT/US01/04789

	85		90		95
Cys Lys Asp Asn Pro Pro Cys Ala Cys Pro Thr Ser Arg Ser Ser Trp					
	100		105		110
Asn Cys Asp Gly Glu Val Leu His Ser Pro Ala Ile Glu Val Arg Val					
	115		120		125
His Cys Gln Leu Val Arg Leu Phe Ala Arg Gly Ile Glu Glu Glu Ser					
	130		135		140

<210> 9
 <211> 382
 <212> DNA
 <213> Mus musculus

<400> 9
 tttatcgagt caagaaattc cacttcacgt cgaagcacgt ggaagacgag gacaatgact 60
 cgaaggaaca agagaagcag aagtttgga agatctgcaa ggacagaccc tcttgcaatt 120
 gctcagcctc cagaagctcc tggaactgag atggcgaagt catgcacagc ccggccattg 180
 aggtcagggt ccactgccag ctggtgcgcc tctttgctcg gggaatcgag gaagagtcac 240
 aagcaagaac cccaaagccc aggagctgtc gcccttgagc tcggggagtg tggaaattac 300
 ttaagaaaaa ttcgacagac cagttatgtt gatatatcca ttggaattta gaaatttatt 360
 tttgataggt aaatcttggt tt 382

<210> 10
 <211> 79
 <212> PRT
 <213> Mus musculus

<400> 10
 Tyr Arg Val Lys Lys Phe His Phe Thr Ser Lys His Val Glu Asp Glu
 1 5 10 15
 Asp Asn Asp Ser Lys Glu Gln Glu Lys Gln Lys Phe Gly Lys Ile Cys
 20 25 30
 Lys Asp Arg Pro Ser Cys Thr Cys Ser Ala Ser Arg Ser Ser Trp Asn
 35 40 45
 Cys Asp Gly Glu Val Met His Ser Pro Ala Ile Glu Val Arg Val His
 50 55 60

WO 01/60990

PCT/US01/04789

Cys Gln Leu Val Arg Leu Phe Ala Arg Gly Ile Glu Glu Glu Ser
65 70 75

<210> 11

<211> 326

<212> PRT

<213> Homo sapiens

<400> 11

Pro Lys His Leu Leu Val Phe Ile Asn Pro Phe Gly Gly Lys Gly Gln
1 5 10 15

Gly Lys Arg Ile Tyr Glu Arg Lys Val Ala Pro Leu Phe Thr Leu Ala
20 25 30

Ser Ile Thr Thr Asp Ile Ile Gly Asn Lys Phe Tyr Val Asn Tyr Val
35 40 45

Glu Val Ile Thr Glu His Ala Asn Gln Ala Lys Glu Thr Leu Tyr Glu
50 55 60

Ile Asn Ile Asp Lys Tyr Asp Gly Ile Val Cys Val Gly Gly Asp Gly
65 70 75 80

Met Phe Ser Glu Val Leu His Gly Leu Ile Gly Arg Thr Gln Arg Ser
85 90 95

Ala Gly Val Asp Gln Asn His Pro Arg Ala Val Leu Val Pro Ser Ser
100 105 110

Leu Arg Ile Gly Ile Ile Pro Ala Gly Ser Thr Asp Cys Val Cys Tyr
115 120 125

Ser Thr Val Gly Thr Ser Asp Ala Glu Thr Ser Ala Leu His Ile Val
130 135 140

Val Gly Asp Ser Leu Ala Met Asp Val Ser Ser Val His His Asn Ser
145 150 155 160

Thr Leu Leu Arg Tyr Ser Val Ser Leu Leu Gly Tyr Gly Phe Tyr Gly
165 170 175

Asp Ile Ile Lys Asp Ser Glu Lys Lys Arg Trp Leu Gly Leu Ala Arg
180 185 190

Tyr Asp Phe Ser Gly Leu Lys Thr Phe Leu Ser His His Cys Tyr Glu
195 200 205

WO 01/60990

PCT/US01/04789

Gly Thr Val Ser Phe Leu Pro Ala Gln His Thr Val Gly Ser Pro Arg
 210 215 220

Asp Arg Lys Pro Cys Arg Ala Gly Cys Phe Val Cys Arg Gln Ser Lys
 225 230 235 240

Gln Gln Leu Glu Glu Glu Gln Lys Lys Ala Leu Tyr Gly Leu Glu Ala
 245 250 255

Ala Glu Asp Val Glu Glu Trp Gln Val Val Cys Gly Lys Phe Leu Ala
 260 265 270

Ile Asn Ala Thr Asn Met Ser Cys Ala Cys Arg Arg Ser Pro Arg Gly
 275 280 285

Leu Ser Pro Ala Ala His Leu Gly Asp Gly Ser Ser Asp Leu Ile Leu
 290 295 300

Ile Arg Lys Cys Ser Arg Phe Asn Phe Leu Arg Phe Leu Ile Arg His
 305 310 315 320

Thr Asn Gln Gln Asp Gln
 325

<210> 12
 <211> 453
 <212> PRT
 <213> Saccharomyces cerevisiae

<400> 12
 Leu Tyr Ile Asp Tyr Lys Pro His Ser Ser Ser His Leu Lys Glu Glu
 1 5 10 15

Asp Asp Leu Val Glu Glu Ile Leu Lys Arg Ser Tyr Lys Asn Thr Arg
 20 25 30

Arg Asn Lys Ser Ile Phe Val Ile Ile Asn Pro Phe Gly Gly Lys Gly
 35 40 45

Lys Ala Lys Lys Leu Phe Met Thr Lys Ala Lys Pro Leu Leu Leu Ala
 50 55 60

Ser Arg Cys Ser Ile Glu Val Val Tyr Thr Lys Tyr Pro Gly His Ala
 65 70 75 80

Ile Glu Ile Ala Arg Glu Met Asp Ile Asp Lys Tyr Asp Thr Ile Ala

WO 01/60990

PCT/US01/04789

85	90	95
Cys Ala Ser Gly Asp Gly Ile Pro His Glu Val Ile Asn Gly Leu Tyr		
100	105	110
Gln Arg Pro Asp His Val Lys Ala Phe Asn Asn Ile Ala Ile Thr Glu		
115	120	125
Ile Pro Cys Gly Ser Gly Asn Ala Met Ser Val Ser Cys His Trp Thr		
130	135	140
Asn Asn Pro Ser Tyr Ser Thr Leu Cys Leu Ile Lys Ser Ile Glu Thr		
145	150	155
Arg Ile Asp Leu Met Cys Cys Ser Gln Pro Ser Tyr Ala Arg Glu His		
165	170	175
Pro Lys Leu Ser Phe Leu Ser Gln Thr Tyr Gly Leu Ile Ala Glu Thr		
180	185	190
Asp Ile Asn Thr Glu Phe Ile Arg Trp Met Gly Pro Ala Arg Phe Glu		
195	200	205
Leu Gly Val Ala Phe Asn Ile Ile Gln Lys Lys Lys Tyr Pro Cys Glu		
210	215	220
Ile Tyr Val Lys Tyr Ala Ala Lys Ser Lys Asn Glu Leu Lys Asn His		
225	230	235
Tyr Leu Glu His Lys Asn Lys Gly Ser Leu Glu Phe Gln His Ile Thr		
245	250	255
Met Asn Lys Asp Asn Glu Asp Cys Asp Asn Tyr Asn Tyr Glu Asn Glu		
260	265	270
Tyr Glu Thr Glu Asn Glu Asp Glu Asp Glu Asp Ala Asp Ala Asp Asp		
275	280	285
Glu Asp Ser His Leu Ile Ser Arg Asp Leu Ala Asp Ser Ser Ala Asp		
290	295	300
Gln Ile Lys Glu Glu Asp Phe Lys Ile Lys Tyr Pro Leu Asp Glu Gly		
305	310	315
Ile Pro Ser Asp Trp Glu Arg Leu Asp Pro Asn Ile Ser Asn Asn Leu		
325	330	335
Gly Ile Phe Tyr Thr Gly Lys Met Pro Tyr Val Ala Ala Asp Thr Lys		

WO 01/60990

PCT/US01/04789

340	345	350
Phe Phe Pro Ala Ala Leu Pro Ser Asp Gly Thr Met Asp Met Val Ile		
355	360	365
Thr Asp Ala Arg Thr Ser Leu Thr Arg Met Ala Pro Ile Leu Leu Gly		
370	375	380
Leu Asp Lys Gly Ser His Val Leu Gln Pro Glu Val Leu His Ser Lys		
385	390	395 400
Ile Leu Ala Tyr Lys Ile Ile Pro Lys Leu Gly Asn Gly Leu Phe Ser		
405	410	415
Val Asp Gly Glu Lys Phe Pro Leu Glu Pro Leu Gln Val Glu Ile Met		
420	425	430
Pro Arg Leu Cys Lys Thr Leu Leu Arg Asn Gly Arg Tyr Val Asp Thr		
435	440	445
Asp Phe Asp Ser Met		
450		

<210> 13
 <211> 436
 <212> PRT
 <213> *Saccharomyces cerevisiae*

<400> 13

Leu Leu Ile Asp His Val Ser Arg Lys Ser Arg Ala Asn Thr Gly Glu
1 5 10 15
Glu Asn Ile Ser Ser Gly Thr Val Glu Glu Ile Leu Glu Lys Ser Tyr
20 25 30
Glu Asn Ser Lys Arg Asn Arg Ser Ile Leu Val Ile Ile Asn Pro His
35 40 45
Gly Gly Lys Gly Thr Ala Lys Asn Leu Phe Leu Thr Lys Ala Arg Pro
50 55 60
Ile Leu Val Glu Ser Gly Cys Lys Ile Glu Ile Ala Tyr Thr Lys Tyr
65 70 75 80
Ala Arg His Ala Ile Asp Ile Ala Lys Asp Leu Asp Ile Ser Lys Tyr
85 90 95

WO 01/60990

PCT/US01/04789

Asp Thr Ile Ala Cys Ala Ser Gly Asp Gly Ile Pro Tyr Glu Val Ile		
100	105	110
Asn Gly Leu Tyr Arg Arg Pro Asp Arg Val Asp Ala Phe Asn Lys Leu		
115	120	125
Ala Val Thr Gln Leu Pro Cys Gly Ser Gly Asn Ala Met Ser Ile Ser		
130	135	140
Cys His Trp Thr Asn Asn Pro Ser Tyr Ala Ala Leu Cys Leu Val Lys		
145	150	155
Ser Ile Glu Thr Arg Ile Asp Leu Met Cys Cys Ser Gln Pro Ser Tyr		
165	170	175
Met Asn Glu Trp Pro Arg Leu Ser Phe Leu Ser Gln Thr Tyr Gly Val		
180	185	190
Ile Ala Glu Ser Asp Ile Asn Thr Glu Phe Ile Arg Trp Met Gly Pro		
195	200	205
Val Arg Phe Asn Leu Gly Val Ala Phe Asn Ile Ile Gln Gly Lys Lys		
210	215	220
Tyr Pro Cys Glu Val Phe Val Lys Tyr Ala Ala Lys Ser Lys Lys Glu		
225	230	235
Leu Lys Val His Phe Leu Glu Asn Lys Asp Lys Asn Lys Gly Cys Leu		
245	250	255
Thr Phe Glu Pro Asn Pro Ser Pro Asn Ser Ser Pro Asp Leu Leu Ser		
260	265	270
Lys Asn Asn Ile Asn Asn Ser Thr Lys Asp Glu Leu Ser Pro Asn Phe		
275	280	285
Leu Asn Glu Asp Asn Phe Lys Leu Lys Tyr Pro Met Thr Glu Pro Val		
290	295	300
Pro Arg Asp Trp Glu Lys Met Asp Ser Glu Leu Thr Asp Asn Leu Thr		
305	310	315
Ile Phe Tyr Thr Gly Lys Met Pro Tyr Ile Ala Lys Asp Thr Lys Phe		
325	330	335
Phe Pro Ala Ala Leu Pro Ala Asp Gly Thr Ile Asp Leu Val Ile Thr		
340	345	350

WO 01/60990

PCT/US01/04789

Asp Ala Arg Ile Pro Val Thr Arg Met Thr Pro Ile Leu Leu Ser Leu
355 360 365

Asp Lys Gly Ser His Val Leu Glu Pro Glu Val Ile His Ser Lys Ile
370 375 380

Leu Ala Tyr Lys Ile Ile Pro Lys Val Glu Ser Gly Leu Phe Ser Val
385 390 395 400

Asp Gly Glu Lys Phe Pro Leu Glu Pro Leu Gln Val Glu Ile Met Pro
405 410 415

Met Leu Cys Lys Thr Leu Leu Arg Asn Gly Arg Tyr Ile Asp Thr Glu
420 425 430

Phe Glu Ser Met
435

<210> 14

<211> 380

<212> PRT

<213> Schizosaccharomyces pombe

<400> 14

Cys Trp Val Asp Phe Val Glu Asn Ser Asp Gln Phe Cys Glu Tyr Leu
1 5 10 15

Leu Asp Val Ala Tyr Lys Gly Ile Lys Arg Ser Arg Arg Phe Ile Val
20 25 30

Phe Ile Asn Pro His Gly Gly Lys Gly Lys Ala Lys His Ile Trp Glu
35 40 45

Ser Glu Ala Glu Pro Val Phe Ser Ser Ala His Ser Ile Cys Glu Val
50 55 60

Val Leu Thr Arg Arg Lys Asp His Ala Lys Ser Ile Ala Lys Asn Leu
65 70 75 80

Asp Val Gly Ser Tyr Asp Gly Ile Leu Ser Val Gly Gly Asp Gly Leu
85 90 95

Phe His Glu Val Ile Asn Gly Leu Gly Glu Arg Asp Asp Tyr Leu Glu
100 105 110

Ala Phe Lys Leu Pro Val Cys Met Ile Pro Gly Gly Ser Gly Asn Ala
115 120 125

WO 01/60990

PCT/US01/04789

Phe	Ser	Tyr	Asn	Ala	Thr	Gly	Gln	Leu	Lys	Pro	Ala	Leu	Thr	Ala	Leu	
130						135					140					
Glu	Ile	Leu	Lys	Gly	Arg	Pro	Thr	Ser	Phe	Asp	Leu	Met	Thr	Phe	Glu	
145					150					155					160	
Gln	Lys	Gly	Lys	Lys	Ala	Tyr	Ser	Phe	Leu	Thr	Ala	Asn	Tyr	Gly	Ile	
				165					170					175		
Ile	Ala	Asp	Cys	Asp	Ile	Gly	Thr	Glu	Asn	Trp	Arg	Phe	Met	Gly	Glu	
			180					185					190			
Asn	Arg	Ala	Tyr	Leu	Gly	Phe	Phe	Leu	Arg	Leu	Phe	Gln	Lys	Pro	Asp	
		195					200					205				
Trp	Lys	Cys	Ser	Ile	Glu	Met	Asp	Val	Val	Ser	Ser	Asp	Arg	Thr	Glu	
	210					215						220				
Ile	Lys	His	Met	Tyr	Glu	Lys	Ser	Lys	Asn	Leu	Ala	Pro	Met	Ser	Glu	
225					230					235					240	
Ser	Ser	Asp	Ser	Asp	Lys	Thr	Val	Ser	Thr	Ser	Pro	Glu	Ser	His	Leu	
				245					250					255		
Leu	Thr	Phe	Glu	Ile	Asn	Asp	Leu	Ser	Ile	Phe	Cys	Ala	Gly	Leu	Leu	
			260					265					270			
Pro	Tyr	Ile	Ala	Pro	Asp	Ala	Lys	Met	Phe	Pro	Ala	Ala	Ser	Asn	Asp	
		275					280					285				
Asp	Gly	Leu	Ile	Asp	Val	Val	Ile	Val	Tyr	Ser	Lys	Gln	Phe	Arg	Lys	
	290					295					300					
Ser	Leu	Leu	Ser	Met	Phe	Thr	Gln	Leu	Asp	Asn	Gly	Gly	Phe	Tyr	Tyr	
305					310					315				320		
Ser	Lys	His	Leu	Asn	Tyr	Tyr	Lys	Val	Arg	Ser	Phe	Arg	Phe	Thr	Pro	
				325					330					335		
Val	Asn	Thr	Gly	Lys	Arg	His	Tyr	Phe	Ala	Leu	Asp	Gly	Glu	Ser	Tyr	
			340					345					350			
Pro	Leu	Glu	Pro	Phe	Glu	Cys	Arg	Val	Ala	Pro	Lys	Leu	Gly	Thr	Thr	
		355					360					365				
Leu	Ser	Pro	Val	Ala	Gly	Phe	Gln	Leu	Leu	Asp	Ile					
		370				375				380						

WO 01/60990

PCT/US01/04789

<210> 15

<211> 415

<212> PRT

<213> *Caenorhabditis elegans*

<400> 15

Cys Arg Ser Asp Ala Glu Glu Asn Glu Gln Leu Thr Ser Val Ile Leu
1 5 10 15

Ser Arg Lys Pro Pro Pro Gln Glu Gln Cys Arg Gly Asn Leu Leu Val
20 25 30

Phe Ile Asn Pro Asn Ser Gly Thr Gly Lys Ser Leu Glu Thr Phe Ala
35 40 45

Asn Thr Val Gly Pro Lys Leu Asp Lys Ser Leu Ile Arg Tyr Glu Val
50 55 60

Val Val Thr Thr Gly Pro Asn His Ala Arg Asn Val Leu Met Thr Lys
65 70 75 80

Ala Asp Leu Gly Lys Phe Asn Gly Val Leu Ile Leu Ser Gly Asp Gly
85 90 95

Leu Val Phe Glu Ala Leu Asn Gly Ile Leu Cys Arg Glu Asp Ala Phe
100 105 110

Arg Ile Phe Pro Thr Leu Pro Ile Gly Ile Val Pro Ser Gly Ser Gly
115 120 125

Asn Gly Leu Leu Cys Ser Val Leu Ser Lys Tyr Gly Thr Lys Met Asn
130 135 140

Glu Lys Ser Val Met Glu Arg Ala Leu Glu Ile Ala Thr Ser Pro Thr
145 150 155 160

Ala Lys Ala Glu Ser Val Ala Leu Tyr Ser Val Lys Thr Asp Asn Gln
165 170 175

Ser Tyr Ala Ser Phe Leu Ser Ile Gly Trp Gly Leu Met Ala Asp Ile
180 185 190

Asp Ile Asp Ser Glu Lys Trp Arg Lys Ser Leu Gly His His Arg Phe
195 200 205

Thr Val Met Gly Phe Ile Arg Ser Cys Asn Leu Arg Ser Tyr Lys Gly

WO 01/60990

PCT/US01/04789

210	215	220
Arg Leu Thr Tyr Arg Pro Tyr Lys Pro Lys Gly Phe His Pro Ser Ser		
225	230	235 240
Asn Val Phe Ser Val Tyr Glu Lys Thr Thr Gln Gln Arg Ile Asp Asp		
	245	250 255
Ser Lys Val Lys Thr Asn Gly Ser Val Ser Asp Ser Glu Glu Glu Thr		
	260	265 270
Met Glu Thr Lys Phe Gln Asn Trp Thr Leu Pro Asp Ser Asp Glu Thr		
	275	280 285
Leu Ala Val Gly Ser Ser Asp Leu Glu Glu Thr Val Val Ile Glu Asp		
	290	295 300
Asn Phe Val Asn Ile Tyr Ala Val Thr Leu Ser His Ile Ala Ala Asp		
	305	310 315 320
Gly Pro Phe Ala Pro Ser Ala Lys Leu Glu Asp Asn Arg Ile His Leu		
	325	330 335
Ser Tyr Ile Leu Trp Lys Asp Ile Gly Thr Arg Val Asn Ile Ala Lys		
	340	345 350
Tyr Leu Leu Ala Ile Glu His Glu Thr His Leu Asp Leu Pro Phe Val		
	355	360 365
Lys His Val Glu Val Ser Ser Met Lys Leu Glu Val Ile Ser Glu Gly		
	370	375 380
Ser His Val Val Leu Asp Gly Glu Val Val Asp Thr Lys Thr Ile Glu		
	385	390 395 400
Val Ala Ser Thr Lys Asn His Ile Ser Val Phe Ser Ser Thr Ala		
	405	410 415

